

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau

(43) International Publication Date  
18 February 2010 (18.02.2010)



(10) International Publication Number  
**WO 2010/019696 A2**

(51) International Patent Classification:  
*C12Q 1/68* (2006.01)

(21) International Application Number:  
PCT/US2009/053589

(22) International Filing Date:  
12 August 2009 (12.08.2009)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
61/088,288 12 August 2008 (12.08.2008) US

(71) Applicant (for all designated States except US): AL-  
LYLIX, INC. [US/US]; 6020 Cornerstone Court, Suite  
260, San Diego, CA 92121 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): JULIEN, Bryan  
[US/US]; 3016 Blackmoor Park Circle, Lexington, KY  
40509 (US). BURLINGAME, Richard [US/US]; 3901  
Rapid Run Drive #828, Lexington, KY 40515 (US).

(74) Agent: LLOYD, Laura, M.; Catalyst Law Group, APC,  
9710 Scranton Road, Suite 170, San Diego, CA 92121  
(US).

(81) Designated States (unless otherwise indicated, for every  
kind of national protection available): AE, AG, AL, AM,  
AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ,  
CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO,  
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,  
HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP,  
KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD,  
ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI,  
NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD,  
SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT,  
TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every  
kind of regional protection available): ARIPO (BW, GH,  
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,  
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ,  
TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,  
MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM,  
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,  
ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished  
upon receipt of that report (Rule 48.2(g))

(54) Title: METHOD FOR PRODUCTION OF ISOPRENOIDS

Figure 1



(57) Abstract: The present invention is directed to variant squalene synthase enzymes, including *Saccharomyces cerevisiae* squalene synthase enzymes, and to nucleic acid molecules encoding these variant enzymes. These variant enzymes produce squalene at a lower rate than the wild-type enzyme, allowing more farnesyl pyrophosphate to be utilized for production of isoprenoid compounds, while still producing sufficient squalene to allow the *S. cerevisiae* cells to grow without the requirement for supplementation by sterols such as ergosterol. These variant enzymes, therefore, are highly suitable for the efficient production of isoprenoids.

WO 2010/019696 A2

## METHOD FOR PRODUCTION OF ISOPRENOIDS

by

Bryan Julien and Richard Burlingame

### CROSS-REFERENCE

[0001] This application claims the benefit of priority to U.S. Provisional Application Serial No. 61/088,288, filed August 12, 2008, which is incorporated herein in its entirety.

### FIELD OF THE INVENTION

[0002] This invention is directed to yeast strains that overproduce a precursor to terpenes, farnesyl pyrophosphate (FPP), and are capable of growing without sterol supplementation.

### BACKGROUND OF THE INVENTION

[0003] With over 50,000 identified members, terpenoids comprise the largest known class of natural products. These compounds are structurally diverse, although based on related carbon skeletons. The structural diversity found among these compounds allows them to perform a variety of essential biochemical functions. These compounds serve as attractants for pollinators, antimicrobial and antiherbivorial defense compounds, and may react with reactive oxygen species to protect against oxidative damage (Dudareva *et al.*, Plant Physiol. 135: 1893-1902 (2004)). As components of the essential oils of aromatic plants, they are largely responsible for the distinct flavors and fragrances associated with their host plants. Moreover, the value of these small molecules extends beyond their biological utility. Many terpenoids have commercial

value as antibiotics, pest control agents, fragrances, flavors, and anti-cancer agents, among other important uses.

[0004] A specific class of these natural products, sesquiterpenoids, is derived from a common 15-carbon building block. This common 15-carbon building block is farnesyl pyrophosphate (FPP). Many important products, such as the flavoring nootkatone, the cosmetic additive bisabolol, and amorpha-4,11-diene, a precursor to the antimalarial compound artemisinin, are sesquiterpenoids and thus are based on the 15-carbon carbon skeleton of FPP. Therefore, methods that can increase the yield of FPP that can be utilized in sesquiterpenoid synthesis are of extreme importance.

[0005] To maximize production of terpenes, mutations in squalene synthase have been used to prevent or minimize conversion of farnesyl pyrophosphate to squalene. In practice, this has been done by either eliminating the corresponding gene, reducing its expression using weak promoters, or controlling its expression with a regulated promoter. Squalene is a precursor to sterols, which are essential to viability of yeast and other organisms. Accordingly, complete elimination of the gene requires feeding of sterols. The yeast *Saccharomyces cerevisiae* is not normally capable of taking up sterols under aerobic conditions, so in order to feed sterols to mutants, secondary mutations enabling sterol uptake are required.

[0006] Various solutions have been proposed in order to obtain high yields of farnesyl pyrophosphate for maximum production of terpenes. In one approach, the *ERG9* gene of the yeast is completely eliminated. The gene *ERG9* encodes the enzyme squalene synthase. However, because these mutants in which *ERG9* is eliminated cannot synthesize squalene, which is a precursor to sterols, they must be fed sterols (Takahashi, *et al.* Biotech. Bioengineer. 97:170-181 (2007)). In another approach, PCT Patent Application Publication No. WO 06/102342 by Bailey *et al.* describes production of high yields of farnesyl pyrophosphate by modifying the expression or activity of one or more polypeptides involved in generating cytosolic acetyl-CoA and/or NADPH. In another approach, a promoter, the *MET3* promoter, is used in place of the native *ERG9* promoter to downregulate the expression of squalene synthase by repressing its synthesis by adding methionine, which acts as a repressor with respect to the *MET3* promoter (Asadollahi *et al.*, Biotech. Bioengineer. 99: 666-677 (2007)). In a similar approach, in addition to

repression of *ERG9* production, overproduction of a soluble, truncated form of 3-hydroxy-3-methylglutaryl-coenzyme A reductase, and enhancement of the activity of the transcription factor *UPC2* was employed (Paradise *et al.*, Cell. Metabol. Engineer. Bioengineer. 100: 371-378 (2008); Ro *et al.*, Nature 440: 940-943 (2006)).

[0007] However, there is a need for improved strains of *Saccharomyces cerevisiae* that can overproduce FPP without the need for sterol supplementation and without regulating expression. Preferably, these improved strains would grow efficiently and produce high levels of farnesyl pyrophosphate for subsequent terpenoid synthesis.

### SUMMARY OF THE INVENTION

[0008] A number of mutations of the *Saccharomyces cerevisiae* squalene synthase gene have been isolated and characterized. These mutants produce a sufficient quantity of squalene synthesis enzyme so that the enzyme catalyzes the synthesis of squalene at a sufficiently high rate so that sterol supplementation for the *S. cerevisiae* cells is not required, while having reduced activity so that more farnesyl pyrophosphate is available for isoprenoid biosynthesis. The reduced activity may be the result of reduced catalytic efficiency of the enzyme, or of reduced intracellular concentration of the protein, or both.

[0009] Accordingly, one aspect of the present invention is an isolated nucleic acid molecule that encodes a squalene synthase enzyme that, when present and expressed *in vivo* in a eukaryotic microbial host as the only squalene synthase species, catalyzes the synthesis of squalene at a sufficiently high rate so that supplementation of the eukaryotic microbial host with a sterol is not required for growth, and also has a reduced squalene synthase activity (referred to herein for convenience as a variant squalene synthase enzyme). In one alternative, a host cell containing the nucleic acid molecule, when expressed *in vivo*, produces a greater concentration of an isoprenoid in grams of isoprenoid per liter of culture than a corresponding host containing a wild-type nucleic acid molecule.

[0010] The variant squalene synthase enzyme of the present invention can be a squalene synthase enzyme of any suitable species. In one aspect, variant squalene synthase enzymes are from *Saccharomyces cerevisiae*.

[0011] Isolated nucleic acid molecules encoding a variant *S. cerevisiae* squalene synthase enzymes according to the present invention include, but are not limited to, the following isolated nucleic acid molecules:

(1) an isolated nucleic acid molecule encoding a variant *S. cerevisiae* squalene synthase enzyme, in which the following nucleic acid changes occur: (a) 691 A→G, resulting in the amino acid change E→G at amino acid residue 149; (b) 748 G→T, resulting in the amino acid change G→V at amino acid residue 168; (c) 786 T→A, resulting in the amino acid change Y→N at amino acid residue 181; (d) 1114 A→T, resulting in the amino acid change Q→L at amino acid residue 290; (e) 1213 T→C, resulting in the amino acid change I→T at amino acid residue 323; and (f) 1290 T→C, resulting in no change of the amino acid L at amino acid residue 349 (silent mutation);

(2) an isolated nucleic acid molecule encoding a variant *S. cerevisiae* squalene synthase enzyme, in which the following nucleic acid changes occur: (a) 72 C→A (in the non-coding region); (b) 110 ΔA (in the non-coding region); and (c) 801 G→A, resulting in the amino acid change V→I at amino acid residue 186;

(3) an isolated nucleic acid molecule encoding a variant *S. cerevisiae* squalene synthase enzyme, in which the following nucleic acid changes occur: (a) 989 T→A, resulting in no change of the amino acid P at amino acid residue 248 (silent mutation); (b) 1112 G→A, resulting in no change of the amino acid E at amino acid residue 289 (silent mutation); (c) 1220 G→A, resulting in no change of the amino acid K at amino acid residue 325 (silent mutation); and (d) 1233 T→C, resulting in the amino acid change Y→H at amino acid residue 330;

(4) an isolated nucleic acid molecule encoding a variant *S. cerevisiae* squalene synthase enzyme, in which the following nucleic acid changes occur: (a) 786 T→A, resulting in the amino acid change Y→N at amino acid residue 181; (b) 1025 A→G, resulting in no change of the amino acid Q at amino acid residue 260 (silent mutation); (c) 1056 T→A, resulting in the amino acid change L→I at amino acid residue

271; (d) 1068 A→G, resulting in the amino acid change S→G at amino acid residue 275; and (e) 1203 A→G, resulting in the amino acid change N→D at amino acid residue 320;

(5) an isolated nucleic acid molecule encoding a variant *S. cerevisiae* squalene synthase enzyme, in which the following nucleic acid changes occur: (a) 886 T→C, resulting in the amino acid change M→T at amino acid residue 214; (b) 969 A→G, resulting in the amino acid change I→V at amino acid residue 242; (c) 1075 T→C, resulting in the amino acid change V→A at amino acid residue 277; and (d) 1114 A→T, resulting in the amino acid change Q→L at amino acid residue 290;

(6) an isolated nucleic acid molecule encoding a variant *S. cerevisiae* squalene synthase enzyme, in which the following nucleic acid changes occur: 84 T→A (in the non-coding region); (b) 283 A→T, resulting in the amino acid change E→V at amino acid residue 13; (c) 424 T→C, resulting in the amino acid change L→P at amino acid residue 60; (d) 440 A→G, resulting in no change of the amino acid R at amino acid residue 65 (silent mutation); and (e) 1076 T→C, resulting in no change of the amino acid V at amino acid residue 277 (silent mutation);

(7) an isolated nucleic acid molecule encoding a variant *S. cerevisiae* squalene synthase enzyme, in which the following nucleic acid changes occur: (a) 619 A→T, resulting in the amino acid change D→V at amino acid residue 125; (b) 634 T→C, resulting in the amino acid change L→P at amino acid residue 130; and (c) 962 C→T, resulting in no change of the amino acid P at amino acid residue 239 (silent mutation);

(8) an isolated nucleic acid molecule encoding a variant *S. cerevisiae* squalene synthase enzyme, in which the following nucleic acid changes occur: (a) 150 A→T (in the non-coding region); (b) 410 T→G, resulting in no change of the amino acid A at amino acid residue 55 (silent mutation); (c) 411 G→T, resulting in the amino acid change V→L at amino acid residue 56; and (d) 1248 T→C, resulting in the amino acid change S→P at amino acid residue 335;

(9) an isolated nucleic acid molecule encoding a variant *S. cerevisiae* squalene synthase enzyme, in which the following nucleic acid changes occur: (a) 510 C→T, resulting in the amino acid change H→Y at amino acid residue 89; (b) 573 T→C, resulting in the amino acid change F→L at amino acid residue 110; (c) 918 A→G,

resulting in the amino acid change R→G at amino acid residue 224; and (d) 997 A→G, resulting in the amino acid change K→G at amino acid residue 251;

(10) an isolated nucleic acid molecule identical to any of (3), (4), (6), (7), or (8), above, except that one or more of the silent mutations in nucleic acid molecules (3), (4), (6), (7), or (8) are omitted;

(11) an isolated nucleic acid molecule encoding a variant *S. cerevisiae* squalene synthase enzyme in which the wild-type *S. cerevisiae* squalene synthase enzyme is mutated with the same amino acid changes as in any of (1) through (10) above;

(12) an isolated nucleic acid molecule encoding a variant *S. cerevisiae* squalene synthase enzyme containing any of the amino acid changes in any of (1) through (10) above; and

(13) an isolated nucleic acid molecule encoding a variant *S. cerevisiae* squalene synthase enzyme in which the squalene synthase enzyme differs from the squalene synthase enzyme encoded by the isolated nucleic acid molecule of any of (1) through (12) above by one to three conservative amino acid substitutions, wherein a conservative amino acid substitution is defined as one of the following substitutions: A→G or S; R→K; N→Q or H; D→E; C→S; Q→N; G→D; G→A or P; H→N or Q; I→L or V; L→I or V; K→R or Q or E; M→L or Y or I; F→M or L or Y; S→T; T→S; W→Y; Y→W or F; and V→I or L.

[0012] The present invention also encompasses an isolated nucleic acid molecule that is at least 95% identical to any of the isolated nucleic acid molecules described above that encodes a variant *S. cerevisiae* squalene synthase enzyme, such that the isolated nucleic acid molecule also encodes a variant *S. cerevisiae* squalene synthase enzyme that, when present and expressed *in vivo* in *Saccharomyces cerevisiae*, catalyzes the synthesis of squalene at a sufficiently high rate that supplementation of the *S. cerevisiae* with a sterol is not required and that has a reduced squalene synthase activity.

[0013] Also within the scope of the present invention is an isolated nucleic acid molecule that includes therein, as a discrete, continuous nucleic acid segment, the isolated nucleic acid molecule encoding the variant squalene synthase as described above. This embodiment of the invention can include, at either the 5'-terminus, the 3'-terminus,

or both, additional nucleic acid sequences such as linkers, adaptors, restriction endonuclease cleavage sites, regulatory sequences such as promoters, enhancers, or operators, or coding sequences, to which the discrete, continuous nucleic acid segment is operatively linked.

[0014] Also within the scope of the present invention are vectors including therein nucleic acid segments according to the present invention as described above, as well as host cells transformed or transfected with the vectors or host cells including therein a nucleic acid segment encoding the variant squalene synthase enzyme according to the present invention, as described above.

[0015] The present invention further includes a variant squalene synthase enzyme encoded by a nucleic acid sequence according to the present invention as described above. The variant squalene synthase enzyme can be, but is not limited to a variant *S. cerevisiae* squalene synthase enzyme. Variant *S. cerevisiae* squalene synthase enzymes according to the present invention include, but are not limited to, at least one of the mutants listed in Table 2.

[0016] Another aspect of the present invention includes a host cell containing and/or expressing a variant squalene synthase enzyme of the present invention as described above (a variant squalene synthase enzyme). The host cell, in this alternative, includes at least one copy of a nucleic acid sequence encoding the enzyme. The copy of the nucleic acid sequence encoding the enzyme can be present in the chromosome of a prokaryotic (bacterial) cell or in one chromosome of a eukaryotic cell. Alternatively, the copy of the nucleic acid sequence encoding the enzyme can be present in a vector or plasmid that is present in the cell.

[0017] Another aspect of the invention is a method of isolating a defective *ERG9* gene. In general, this method comprises the steps of:

- (1) isolating a wild-type *ERG9* gene to produce an isolated wild-type *ERG9* gene;
- (2) subjecting the isolated wild-type *ERG9* gene to mutagenesis to generate a pool of *erg9* mutants;
- (3) transforming mutants from the pool of *erg9* mutants generated in step (2) into a strain of a eukaryotic microbial host that contains a plasmid expressing a



terpene synthase gene that produces a detectable and measurable terpene product, the strain of the eukaryotic microbial host being transformed in such a manner that replacement of the preexisting *ERG9* allele with an *erg9* mutation allows the strain to grow in a sterol-free medium; and

(4) isolating a transformant from step (3) that produces a level of terpene product at least equivalent to the level of terpene product produced by a strain of the eukaryotic microbial host expressing the terpene synthase gene that requires a sterol in the medium for growth.

[0018] Another aspect of the present invention is a method of isolating a variant squalene synthase enzyme. The variant squalene synthase enzyme to be isolated by the methods of the invention is as described above.

[0019] In general, this method comprises the steps of:

- (a) culturing a host cell that expresses a variant squalene synthase gene according to the present invention or that contains a variant squalene synthase enzyme according to the present invention; and
- (b) isolating the variant squalene synthase enzyme from the host cell.

[0020] Yet another aspect of the present invention is a method of producing an isoprenoid using a host cell containing a mutated *ERG9* gene, which defective *ERG9* gene encodes a variant squalene synthase enzyme.

[0021] In one alternative, a host cell that includes a mutated *ERG9* gene encoding a variant squalene synthase enzyme further includes at least one isoprenoid synthase gene, so that the farnesyl pyrophosphate produced in the host cell, which is available in greater concentrations for isoprenoid biosynthesis can be converted to an isoprenoid by the isoprenoid synthase encoded by the isoprenoid synthase gene.

[0022] This alternative, in general, comprises the steps of:

- (1) providing a host cell that includes a mutated *ERG9* gene that encodes a variant squalene synthase enzyme according to the present invention and at least one isoprenoid synthase gene;
- (2) allowing the host cell to produce farnesyl pyrophosphate and to synthesize the isoprenoid from the farnesyl pyrophosphate; and
- (3) isolating the isoprenoid synthesized by the host cell.

[0023] In another alternative method for producing an isoprenoid, the method, in general, comprises the steps of:

- (1) providing a host cell that includes a mutated *ERG9* gene that encodes a variant synthase enzyme according to the present invention;
- (2) allowing the host cell to produce farnesyl pyrophosphate;
- (3) isolating farnesyl pyrophosphate from the host cell;
- (4) reacting the farnesyl pyrophosphate *in vitro* with one or more isoprenoid synthases to synthesize the isoprenoid; and
- (5) isolating the isoprenoid.

### BRIEF DESCRIPTION OF THE DRAWINGS

[0024] These and other features, aspects, and advantages of the present invention will become better understood with regard to the following description, appended claims, and accompanying drawings.

[0025] Figure 1 depicts a graph showing the concentration of premnaspirodiene in mg/L for each of the 25 isolates grown with and without ergosterol supplementation.

[0026] Figure 2 depicts shows the sequences of the wild-type *ERG9* gene including sequences 245 base pairs upstream of the start site (SEQ ID NO: 3). In Figure 2, the underlined nucleotides shown at the 5'-terminus and 3'-terminus of the *ERG9* gene sequence represent the upstream primer (7-162.1) 5'-CCATCTTCAACAACAATACCG-3' (SEQ ID NO: 1) and the downstream primer (7-162.2) 5'-GTACTTAGTTATTGTTTCGG-3' (SEQ ID NO: 2).

### DETAILED DESCRIPTION OF THE INVENTION

#### Definitions

[0027] As used herein, the term “nucleic acid,” “nucleic acid sequence,” “polynucleotide,” or similar terms, refers to a deoxyribonucleotide or ribonucleotide oligonucleotide or polynucleotide, including single- or double-stranded forms, and coding or non-coding (*e.g.*, “antisense”) forms. The term encompasses nucleic acids containing known analogues of natural nucleotides. The term also encompasses nucleic acids including modified or substituted bases as long as the modified or substituted bases interfere neither with the Watson-Crick binding of complementary nucleotides or with the binding of the nucleotide sequence by proteins that bind specifically. The term also encompasses nucleic-acid-like structures with synthetic backbones. DNA backbone analogues provided by the invention include phosphodiester, phosphorothioate, phosphorodithioate, methylphosphonate, phosphoramidate, alkyl phosphotriester, sulfamate, 3'-thioacetal, methylene(methylimino), 3'-N-carbamate, morpholino carbamate, and peptide nucleic acids (PNAs). (Oligonucleotides and Analogues, a Practical Approach, edited by F. Eckstein, IRL Press at Oxford University Press (1991);

Antisense Strategies, Annals of the New York Academy of Sciences, Volume 600, Eds. Baserga and Denhardt (NYAS 1992); Milligan, J. Med. Chem. 36:1923-1937 (1993); Antisense Research and Applications (1993, CRC Press). PNAs contain non-ionic backbones, such as N-(2-aminoethyl) glycine units. Phosphorothioate linkages are described in, *e.g.* U.S. Pat. Nos. 6,031,092; 6,001,982; 5,684,148; WO 97/03211; WO 96/39154; Mata, Toxicol. Appl. Pharmacol. 144:189-197 (1997). Other synthetic backbones encompassed by the term include methylphosphonate linkages or alternating methylphosphonate and phosphodiester linkages (U.S. Pat. No. 5,962,674; Strauss-Soukup, Biochemistry 36:8692-8698 (1997)), and benzylphosphonate linkages (U.S. Pat. No. 5,532,226; Samstag, Antisense Nucleic Acid Drug Dev 6:153-156 (1996)). Bases included in nucleic acids include any of the known base analogs of DNA and RNA including, but not limited to, 4-acetylcytosine, 8-hydroxy-N<sup>6</sup>-methyladenosine, aziridinylcytosine, pseudoisocytosine, 5-(carboxyhydroxymethyl)uracil, 5-fluorouracil, 5-bromouracil, 5-carboxymethylaminomethyl-2-thiouracil, 5-carboxymethylaminomethyluracil, dihydrouracil, inosine, N<sup>6</sup>-isopentenyladenine, 1-methyladenine, 1-methylpseudo-uracil, 1-methylguanine, 1-methylinosine, 2,2-dimethyl-guanine, 2-methyladenine, 2-methylguanine, 3-methyl-cytosine, 5-methylcytosine, N<sup>6</sup>-methyladenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxy-amino-methyl-2-thiouracil,  $\beta$ -D-mannosylqueosine, 5'-methoxycarbonylmethyluracil, 5-methoxyuracil, 2-methylthio-N<sup>6</sup>-isopentenyladenine, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, oxybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, N-uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, pseudouracil, queosine, 2-thiocytosine, and 2,6-diaminopurine. DNA may be in the form of cDNA, in vitro polymerized DNA, plasmid DNA, parts of a plasmid DNA, genetic material derived from a virus, linear DNA, vectors (*e.g.* P1, PAC, BAC, YAC, and artificial chromosomes), expression cassettes, chimeric sequences, recombinant DNA, chromosomal DNA, an oligonucleotide, anti-sense DNA, or derivatives of these groups. RNA may be in the form of oligonucleotide RNA, tRNA (transfer RNA), snRNA (small nuclear RNA), rRNA (ribosomal RNA), mRNA (messenger RNA), *in vitro* polymerized RNA, recombinant RNA, chimeric sequences, anti-sense RNA, siRNA (small interfering RNA), ribozymes, or derivatives of

these groups. Additionally, the terms “nucleic acid” or “nucleic acid molecule” refer to a deoxyribonucleotide or ribonucleotide polymer in either single-or double-stranded form, and unless otherwise limited, would encompass known analogs of natural nucleotides that can function in a similar manner as naturally occurring nucleotides. A “nucleotide sequence” also refers to a polynucleotide molecule or oligonucleotide molecule in the form of a separate fragment or as a component of a larger nucleic acid. The nucleotide sequence or molecule may also be referred to as a “nucleotide probe.” Some of the nucleic acid molecules of the invention are derived from DNA or RNA isolated at least once in substantially pure form and in a quantity or concentration enabling identification, manipulation, and recovery of its component nucleotide sequence by standard biochemical methods. Examples of such methods, including methods for PCR protocols that may be used herein, are disclosed in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd ed., Cold Spring Harbor Laboratory Press, New York (1989), Ausubel, F. A., et al., eds., *Current Protocols in Molecular Biology*, John Wiley and Sons, Inc., New York (1987), and Innis, M., et al. (Eds.) *PCR Protocols: A Guide to Methods and Applications*, Academic Press, San Diego, Calif. (1990). Reference to a nucleic acid molecule also includes its complement as determined by the standard Watson-Crick base-pairing rules, with uracil (U) in RNA replacing thymine (T) in DNA where necessary, unless the complement is specifically excluded.

[0028] As described herein, the nucleic acid molecules of the invention include DNA in both single-stranded and double-stranded form, as well as the DNA or RNA complement thereof. DNA includes, for example, DNA, genomic DNA, chemically synthesized DNA, DNA amplified by PCR, and combinations thereof. Genomic DNA, including translated, non-translated and control regions, may be isolated by conventional techniques, *e.g.*, using any one of the cDNAs of the invention, or suitable fragments thereof, as a probe, to identify a piece of genomic DNA which can then be cloned using methods commonly known in the art.

[0029] Polypeptides encoded by the nucleic acids of the invention are encompassed by the invention. As used herein, reference to a nucleic acid “encoding” a protein or polypeptide encompasses not only cDNAs and other intronless nucleic acids, but also DNAs, such as genomic DNA, with introns, on the assumption that the introns

included have appropriate splice donor and acceptor sites that will ensure that the introns are spliced out of the corresponding transcript when the transcript is processed in a eukaryotic cell. Due to the degeneracy of the genetic code wherein more than one codon can encode the same amino acid, multiple DNA sequences can code for the same polypeptide. Such variant DNA sequences can result from genetic drift or artificial manipulation (*e.g.*, occurring during PCR amplification or as the product of deliberate mutagenesis of a native sequence). Deliberate mutagenesis of a native sequence can be carried out using numerous techniques well known in the art. For example, oligonucleotide-directed site-specific mutagenesis procedures can be employed, particularly where it is desired to mutate a gene such that predetermined restriction nucleotides or codons are altered by substitution, deletion or insertion. Exemplary methods of making such alterations are disclosed by Walder *et al.*, *Gene*, 42:133 (1986); Bauer *et al.*, *Gene* 37:73 (1985); Craik, *BioTechniques*, Jan. 12-19 (1985); Smith *et al.*, *Genetic Engineering: Principles and Methods*, Plenum Press, (1981); Kunkel (PNAS USA 82:488 (1985); Kunkel *et al.*, *Methods in Enzymol.* 154:367 (1987). The present invention thus encompasses any nucleic acid capable of encoding a protein of the current invention.

[0030] In a peptide or protein, suitable conservative substitutions of amino acids are known to those of skill in this art and may be made generally without altering the biological activity of the resulting molecule. Those of skill in this art recognize that, in general, single amino acid substitutions in non-essential regions of a polypeptide do not substantially alter biological activity (see, *e.g.* Watson *et al.*, *Molecular Biology of the Gene*, 4th Edition, 1987, Benjamin/Cummings, p. 224). In particular, such a conservative variant has a modified amino acid sequence, such that the change(s) do not substantially alter the proteins structure and/or activity, *e.g.*, antibody activity, enzymatic activity, or receptor activity. These include conservatively modified variations of an amino acid sequence, *i.e.*, amino acid substitutions, additions or deletions of those residues that are not critical for protein activity, or substitution of amino acids with residues having similar properties (*e.g.*, acidic, basic, positively or negatively charged, polar or non-polar, etc.) such that the substitutions of even critical amino acids does not substantially alter structure and/or activity. Conservative substitution tables providing functionally similar

amino acids are well known in the art. For example, one exemplary guideline to select conservative substitutions includes (original residue followed by exemplary substitution): Ala/Gly or Ser; Arg/Lys; Asn/Gln or His; Asp/Glu; Cys/Ser; Gln/Asn; Gly/Asp; Gly/Ala or Pro; His/Asn or Gln; Ile/Leu or Val; Leu/Ile or Val; Lys/Arg or Gln or Glu; Met/Leu or Tyr or Ile; Phe/Met or Leu or Tyr; Ser/Thr; Thr/Ser; Trp/Tyr; Tyr/Trp or Phe; Val/Ile or Leu. An alternative exemplary guideline uses the following six groups, each containing amino acids that are conservative substitutions for one another: (1) alanine (A or Ala), serine (S or Ser), threonine (T or Thr); (2) aspartic acid (D or Asp), glutamic acid (E or Glu); (3) asparagine (N or Asn), glutamine (Q or Gln); (4) arginine (R or Arg), lysine (K or Lys); (5) isoleucine (I or Ile), leucine (L or Leu), methionine (M or Met), valine (V or Val); and (6) phenylalanine (F or Phe), tyrosine (Y or Tyr), tryptophan (W or Trp); (Creighton (1984) *Proteins*, W. H. Freeman and Company; Schulz and Schimer (1979) *Principles of Protein Structure*, Springer-Verlag). One of skill in the art will appreciate that the above-identified substitutions are not the only possible conservative substitutions. For example, for some purposes, one may regard all charged amino acids as conservative substitutions for each other whether they are positive or negative. In addition, individual substitutions, deletions or additions that alter, add or delete a single amino acid or a small percentage of amino acids in an encoded sequence can also be considered "conservatively modified variations" when the three-dimensional structure and the function of the protein to be delivered are conserved by such a variation.

[0031] As used herein, the term "isolated" with reference to a nucleic acid molecule or polypeptide or other biomolecule means that the nucleic acid or polypeptide has been separated from the natural environment from which the polypeptide or nucleic acid were obtained. It may also mean that the biomolecule has been altered from the natural state. For example, a polynucleotide or a polypeptide naturally present in a living animal is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated," as the term is employed herein. Thus, a polypeptide or polynucleotide produced and/or contained within a recombinant host cell is considered isolated. Also intended as an "isolated polypeptide" or an "isolated polynucleotide" are polypeptides or polynucleotides that have been purified, partially or substantially, from a recombinant host cell or from a native source. For

example, a recombinantly produced version of a compound can be substantially purified by the one-step method described in Smith *et al.*, Gene 67:3140 (1998). The terms “isolated” and “purified” are sometimes used interchangeably.

[0032] Thus, by “isolated” it is meant that the nucleic acid is free of the coding sequences of those genes that, in a naturally-occurring genome, immediately flank the gene encoding the nucleic acid of interest. Isolated DNA may be single-stranded or double-stranded, and may be genomic DNA, cDNA, recombinant hybrid DNA, or synthetic DNA. It may be identical to a native DNA sequence, or may differ from such sequence by the deletion, addition, or substitution of one or more nucleotides.

[0033] “Isolated” or “purified” also refer to preparations made from biological cells or hosts means any cell extract containing the indicated DNA or protein including a crude extract of the DNA or protein of interest. For example, in the case of a protein, a purified preparation can be obtained following an individual technique or a series of preparative or biochemical techniques and the protein of interest can be present at various degrees of purity in these preparations. Particularly for proteins, the procedures may include for example, but are not limited to, ammonium sulfate fractionation, gel filtration, ion exchange change chromatography, affinity chromatography, density gradient centrifugation, electrofocusing, chromatofocusing, and electrophoresis. As used herein, the term “substantially purified,” when applied to a composition or extract derived from yeast, and wherein the composition or extract contains an isoprenoid, is hereby defined as containing at least about twice the concentration of isoprenoid in proportion to yeast material, wherein yeast material is defined as being selected from the group consisting of yeast cell membrane, yeast organelle, yeast cytoplasm, yeast microsomal fraction, yeast cell, and yeast extract.

[0034] A preparation of DNA or protein that is “substantially pure” or “isolated” refers to a preparation free from naturally occurring materials with which such DNA or protein is normally associated in nature. “Essentially pure” means a “highly” purified preparation that contains at least 95% of the DNA or protein of interest.

[0035] A cell extract that contains the DNA or protein of interest should be understood to mean a homogenate preparation or cell-free preparation obtained from cells



that express the protein or contain the DNA of interest. The term "cell extract includes culture media, for example, spent culture media from which the cells have been removed.

[0036] A "vector" is a nucleic acid that is capable of transporting another nucleic acid. Vectors may be, for example, plasmids, viruses, cosmids or phage. An "expression vector" is a vector that is capable of directing expression of a protein encoded by one or more genes carried by the vector when it is present in the appropriate environment. Examples of vectors are those that can autonomously replicate and express structural gene products present in the DNA segments to which they are operatively linked. Vectors, therefore, can contain the replicons and selectable markers described earlier. Vectors include, but are not necessarily limited to, expression vectors.

[0037] As used herein with regard to nucleic acid molecules, including DNA fragments, the phrase "operatively linked" means the sequences or segments have been covalently joined, preferably by conventional phosphodiester bonds, into one strand of DNA, whether in single or double-stranded form such that operatively linked portions function as intended.

[0038] As used herein, the phrase "substantially identical" means that a relevant sequence is at least 70%, 75%, 80%, 85%, 90%, 92%, 95%, 96%, 97%, 98%, or 99% identical to a given sequence. By way of example, such sequences may be allelic variants, sequences derived from various species, or they may be derived from the given sequence by truncation, deletion, amino acid substitution or addition. Percent identity between two sequences is determined by standard alignment algorithms such as ClustalX when the two sequences are in best alignment according to the alignment algorithm.

[0039] In order to maximize production of farnesyl pyrophosphate (FPP), novel mutants in the *ERG9* gene of *Saccharomyces cerevisiae* have been developed. The *ERG9* gene is the gene encoding squalene synthase. These mutants have reduced, but not eliminated, squalene synthase activity. As such, they allow sufficient production of squalene and subsequent sterols to allow growth, but are sufficiently reduced in activity to allow accumulation of FPP and overproduction of terpenes. This is done, unlike in previous approaches, by generating and utilizing defective squalene synthase genes, which when expressed result in reduced cellular squalene synthase activity rather than

downregulating the transcription of a normally active squalene synthase enzyme. This makes the reduced squalene synthase activity independent of the activity of a repressor.

[0040] Accordingly, one aspect of the present invention is an isolated nucleic acid molecule that encodes a squalene synthase enzyme that, when present and expressed *in vivo* in a eukaryotic microbial host cell, catalyzes the synthesis of squalene at a sufficiently high rate that supplementation of the eukaryotic microbial host cell with a sterol is not required and that has a reduced squalene synthase activity (referred to herein for convenience as a variant squalene synthase enzyme). Typically, the variant squalene synthase enzyme encoded by the isolated nucleic acid molecule of the present invention has a reduced  $V_{\max}$  for squalene synthesis.  $V_{\max}$  is the maximum rate of a reaction being catalysed by an enzyme. Alternatively, the variant squalene synthase enzyme encoded by the isolated nucleic acid molecule of the present invention has an increased Michaelis constant ( $K_m$ ) for its FPP substrate, in which case the enzyme is less active at a given intracellular concentration of FPP than the wild-type enzyme. The  $K_m$  is a means of characterising an enzyme's affinity for a substrate. The  $K_m$  in an enzymatic reaction is the substrate concentration at which the reaction rate is half its maximum speed. Typically, the squalene synthase enzyme encoded by the isolated nucleic acid molecule of the present invention, when expressed *in vivo* in the eukaryotic microbial host cell, produces squalene at a rate of less than 75% of the wild-type enzyme. Preferably, the squalene synthase enzyme encoded by the isolated nucleic acid molecule of the present invention, when expressed *in vivo* in the eukaryotic microbial host cell, produces squalene at a rate of less than 50% of the wild-type enzyme. More preferably, the squalene synthase enzyme encoded by the isolated nucleic acid molecule of the present invention, when expressed *in vivo* in the eukaryotic microbial host cell, produces squalene at a rate of less than 25% of the wild-type enzyme. The eukaryotic microbial host cell is typically, but is not limited to, a fungal host cell. The fungal host cell is typically, but it not limited to, a yeast host cells, such as a *Saccharomyces cerevisiae* host cell or other host cells of the genus *Saccharomyces*. Similarly, the variant squalene synthase is not limited to a squalene synthase of *Saccharomyces cerevisiae*, but can be a squalene synthase of another species of *Saccharomyces* or a squalene synthase of any

organism that has a gene that catalyzes the conversion of farnesyl pyrophosphate into squalene.

[0041] Isolated nucleic acid molecules according to the present invention include, but are not limited to, the following isolated nucleic acid molecules:

- (1) an isolated nucleic acid molecule encoding a variant *S. cerevisiae* squalene synthase enzyme as shown as Mutant 4 in Table 2 of the present invention;
- (2) an isolated nucleic acid molecule encoding a mutated *S. cerevisiae* squalene synthase enzyme as shown as Mutant 10 in Table 2 of the present invention;
- (3) an isolated nucleic acid molecule encoding a mutated *S. cerevisiae* squalene synthase enzyme as shown as Mutant 14 in Table 2 of the present invention;
- (4) an isolated nucleic acid molecule encoding a mutated *S. cerevisiae* squalene synthase enzyme as shown as Mutant 19 in Table 2 of the present invention;
- (5) an isolated nucleic acid molecule encoding a mutated *S. cerevisiae* squalene synthase enzyme as shown as Mutant 22 in Table 2 of the present invention;
- (6) an isolated nucleic acid molecule encoding a mutated *S. cerevisiae* squalene synthase enzyme as shown as Mutant 23 in Table 2 of the present invention;
- (7) an isolated nucleic acid molecule encoding a mutated *S. cerevisiae* squalene synthase enzyme as shown as Mutant 24 in Table 2 of the present invention;
- (8) an isolated nucleic acid molecule encoding a mutated *S. cerevisiae* squalene synthase enzyme as shown as Mutant 25 in Table 2 of the present invention;
- (9) an isolated nucleic acid molecule encoding a mutated *S. cerevisiae* squalene synthase enzyme as shown as Mutant 69 in Table 2 of the present invention;
- (10) an isolated nucleic acid molecule identical to any of (3), (4), (6), (7), or (8), above, except that one or more of the silent mutations in nucleic acid molecules (3), (4), (6), (7), or (8) are omitted;
- (11) an isolated nucleic acid molecule encoding a variant *S. cerevisiae* squalene synthase protein in which the wild-type *S. cerevisiae* squalene synthase enzyme is mutated with the same amino acid changes as in any of (1) through (10) above;
- (12) an isolated nucleic acid encoding a squalene synthase protein containing any of the amino acid changes in any of (1) through (10). Although some of the mutations described above are designated as "silent," meaning that they do not affect

the amino acid inserted into the polypeptide chain at that position, there is evidence that such mutations may affect protein function in a number of ways, including altering folding patterns due to effects on translation due to codon utilization.

[0042] The invention includes an isolated nucleic acid molecule encoding a squalene synthase enzyme that differs from the squalene synthase enzyme encoded by the nucleic acid molecule in any of (1) through (11), above, by one to three conservative amino acid substitutions, in which a conservative amino acid substitution is defined as one of the following substitutions: A→G or S; R→K; N→Q or H; D→E; C→S; Q→N; G→D; G→A or P; H→N or Q; I→L or V; L→I or V; K→R or Q or E; M→L or Y or I; F→M or L or Y; S→T; T→S; ;W→Y; Y→W or F; and V→I or L. Preferably, the isolated nucleic acid molecule encodes a squalene synthase protein that differs from the squalene synthase protein encoded by the nucleic acid molecule in any of (1) through (10) by one or two conservative amino acid substitutions. More preferably, the isolated nucleic acid molecule encodes a squalene synthase protein that differs from the squalene synthase protein encoded by the nucleic acid molecule in any of (1) through (11) by one conservative amino acid substitution.

[0043] The invention includes an isolated nucleic acid molecule that is at least 95% identical to any of the isolated nucleic acid molecules described above that encodes a mutated *S. cerevisiae* squalene synthase enzyme, such that the isolated nucleic acid molecule also encodes a mutated *S. cerevisiae* squalene synthase enzyme that, when present and expressed *in vivo* in *Saccharomyces cerevisiae*, catalyzes the synthesis of squalene at a sufficiently high rate that supplementation of the *S. cerevisiae* with sterols is not required and that has a reduced squalene synthase activity. Typically, the isolated nucleic acid molecule is at least 97.5% identical to any of the isolated nucleic acid molecules described above. Preferably, the isolated nucleic acid molecule is at least 99% identical to any of the isolated nucleic acid molecules described above. More preferably, the isolated nucleic acid molecule is at least 99.5% identical to any of the isolated nucleic acid molecules described above. Most preferably, the isolated nucleic acid molecule is at least 99.8% identical to any of the isolated nucleic acid molecules described above. For these purposes, "identity" is defined according to the Needleman-Wunsch algorithm (S.B. Needleman & C.D. Wunsch, "A General Method Applicable to the Search for

Similarities in the Amino Acid Sequence of Two Proteins," J. Mol. Biol. 48: 443-453 (1970)).<sup>6</sup>

[0044] Nucleic acid molecules according to the present invention and having the desired degree of identity are not limited to nucleic acid molecules derived from *S. cerevisiae*; they include nucleic acid molecules derived from other species of *Saccharomyces*, or, as described above, derived from any organism that has a gene capable of catalyzing the conversion of farnesyl pyrophosphate into squalene.

[0045] Additionally, isolated nucleic acid molecules according to the present invention further include isolated nucleic acid molecules, which encode a squalene synthase enzyme, and when expressed in a eukaryotic microbial host in which no other squalene synthase enzyme is expressed, result in a significant reduction of conversion of farnesyl pyrophosphate to squalene as described above. In one alternative, a nucleic acid molecule according to the present invention, when expressed *in vivo*, causes a host cell to produce a greater concentration of an isoprenoid in grams of isoprenoid per liter of culture than a corresponding host cell expressing a wild-type nucleic acid molecule. These isolated nucleic acid molecules have at least one change from a nucleic acid molecule that includes the coding region for the wild-type *ERG9* gene and its flanking sequences, including the sequences both upstream and downstream from the coding region. This at least one change reduces the squalene synthase activity, even though the specific activity may potentially be unaltered. The reduction of the activity of the squalene synthase enzyme can occur through one or more of the following mechanisms: (1) reduction in transcription so that less mRNA that can be translated into squalene synthase enzyme is generated; (2) reduction of mRNA stability, again reducing translation; and (3) reduction of enzyme stability brought about by an increased rate of protein degeneration *in vivo*. In other words, either: (1) the specific activity of the resulting squalene synthase enzyme is reduced through at least one change in the amino acid sequence of the enzyme expressed from the nucleic acid molecule; or (2) the *in vivo* activity of the enzyme is reduced through a reduction in transcription, a reduction in translation, or a reduction of enzyme stability.

[0046] The nucleic acid described above can be DNA, RNA, or a RNA-DNA hybrid, but is typically DNA. The nucleic acid described above can be single-stranded or

double-stranded. If the nucleic acid is single-stranded, either the strand described or its complement can be the coding strand and is within the scope of the invention.

[0047] Also within the scope of the invention is an isolated nucleic acid molecule that includes therein, as a discrete, continuous nucleic acid segment, the isolated nucleic acid molecule encoding the variant squalene synthase. This embodiment of the invention can include, at either the 5'-terminus, the 3'-terminus, or both, additional nucleic acid sequences such as linkers, adaptors, restriction endonuclease cleavage sites, regulatory sequences such as promoters, enhancers, or operators, or coding sequences, to which the discrete, continuous nucleic acid segment is operatively linked. In the event that the isolated nucleic acid molecule includes additional coding sequences, the isolated nucleic acid molecule can encode a fusion protein having *S. cerevisiae* squalene synthase activity.

[0048] Also within the scope of the invention are vectors including therein nucleic acid segments according to the present invention as described above. The vectors can be capable of replication in prokaryotes (bacteria) or in eukaryotes (yeast or cells of higher organisms). In one alternative, the vectors are capable of replication in yeast, for example, *S. cerevisiae*. In yeast, a number of vectors containing constitutive or inducible promoters may be used. For a review, see Current Protocols in Molecular Biology, Vol. 2, 1988, Ed. Ausubel, et al., Greene Publish. Assoc. & Wiley Interscience, Ch. 13; Grant, et al., 1987, Expression and Secretion Vectors for Yeast, in Methods in Enzymology, Eds. Wu & Grossman, 31987, Acad. Press, N.Y., Vol. 153, pp.516-544; Glover, 1986, DNA Cloning, Vol. II, IRL Press, Wash., D.C., Ch. 3; and Bitter, 1987, Heterologous Gene Expression in Yeast, Methods in Enzymology, Eds. Berger & Kimmel, Acad. Press, N.Y., Vol. 152, pp. 673-684; and The Molecular Biology of the Yeast *Saccharomyces*, 1982, Eds. Strathern et al., Cold Spring Harbor Press, Vols. I and II. A constitutive yeast promoter such as ADH1 or LEU2 or an inducible promoter such as GAL4 may be used (Cloning in Yeast, Ch. 3, Rothstein In: DNA Cloning Vol. II, A Practical Approach, Ed. DM Glover, 1986, IRL Press, Wash., D.C.). Alternatively, vectors may be used which promote integration of foreign DNA sequences into the yeast chromosome.

[0049] Appropriate cloning and expression vectors for use with bacterial, fungal, yeast, and mammalian cellular hosts are described, for example, in Pouwels et al.,

Cloning Vectors. A Laboratory Manual, Elsevier, N.Y., (1985). Cell-free translation systems could also be employed to produce the disclosed polypeptides using RNAs derived from DNA constructs disclosed herein.

[0050] Examples of expression vectors that can be used in prokaryotic host cells include those derived from commercially available plasmids such as the cloning vector pET plasmids (Novagen, Madison, Wis., USA) or pBR322 (ATCC 37017). The pBR322 vector contains genes for ampicillin and tetracycline resistance and thus provides simple means for identifying transformed cells. To construct an expression vector using pBR322, an appropriate promoter and a DNA sequence encoding one or more of the polypeptides of the invention are inserted into the pBR322 vector. Other commercially available vectors include, for example, pKK223-3 (Pharmacia Fine Chemicals, Uppsala, Sweden) and pGEM-1 (Promega Biotec, Madison, Wis., USA). Other commercially available vectors include those that are specifically designed for the expression of proteins; these would include pMAL-p2 and pMAL-c2 vectors that are used for the expression of proteins fused to maltose binding protein (New England Biolabs, Beverly, Mass., USA).

[0051] Promoter sequences commonly used for recombinant prokaryotic host cell expression vectors include the bacteriophage T7 promoter (Studier and Moffatt, J. Mol. Biol. 189:113 (1986)),  $\beta$ -lactamase (penicillinase), lactose promoter system (Chang *et al.*, Nature 275:615, 1978; Goeddel *et al.*, Nature 281:544 (1979)), tryptophan (trp) promoter system (Goeddel *et al.*, Nucl. Acids Res. 8:4057 (1980); EP-A-36776), and tac promoter (Maniatis, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, p. 412(1982)). A particularly useful prokaryotic host cell expression system employs a phage  $\lambda$  PL promoter and a c1857ts thermolabile repressor sequence. Plasmid vectors available from the American Type Culture Collection (ATCC), which incorporate derivatives of the  $P_L$  promoter, include plasmid pHUB2 (resident in *E. coli* strain JMB9 (ATCC 37092)) and pPLc28 (resident in *E. coli* RR1 (ATCC 53082)).

[0052] As detailed below, nucleic acid segments according to the present invention can also be incorporated in vectors suitable for introduction into yeast cells, such as, for example, *Saccharomyces* (particularly *S. cerevisiae*), *Pichia* (particularly *P. pastoris*), and *Kluyveromyces* (particularly *K. lactis*). Yeast vectors will often contain an

origin of replication sequence from a 2 $\mu$  yeast plasmid, an autonomously replicating sequence (ARS), a promoter region, sequences for polyadenylation, sequences for transcription termination, and a selectable marker gene. Suitable promoter sequences for yeast vectors include, among others, promoters for metallothionein, 3-phosphoglycerate kinase (Hitzeman *et al.*, J. Biol. Chem. 255:2073, 1980), or other glycolytic enzymes (Hess *et al.*, J. Adv. Enzyme Reg. 7:149 (1968); Holland *et al.*, Biochem. 17:4900, (1978)), such as enolase, glyceraldehyde phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase. Other suitable vectors and promoters for use in yeast expression are described in Hitzeman, EPA-73,657 or in Fleer *et al.*, Gene, 107:285-195 (1991); and van den Berg *et al.*, Bio/Technology, 8:135-139 (1990). Another alternative is the glucose-repressible ADH2 promoter described by Russell *et al.* (J. Biol. Chem. 258:2674, (1982)) and Beier *et al.* (Nature 300:724 (1982)). Shuttle vectors replicable in both yeast and *E. coli* can be constructed by inserting DNA sequences from pBR322 for selection and replication in *E. coli* (Amp<sup>r</sup> gene and origin of replication) into the above-described yeast vectors.

[0053] When the vectors are capable of replication in cells of higher organisms, the higher organisms can be plants or animals, including mammals. In cases where plant expression vectors are used, the expression of a mutated *S. cerevisiae* squalene synthase coding sequence according to the present invention may be driven by any of a number of promoters. For example, viral promoters such as the 35S RNA and 19S RNA promoters of CaMV (Brisson *et al.*, Nature, 310:511-514 (1984)), or the coat protein promoter to TMV (Takamatsu *et al.*, EMBO J., 6:307-311 (1987)) may be used; alternatively, plant promoters such as the small subunit of RUBISCO (Coruzzi *et al.*, EMBO J. 3:1671-1680 (1984); Broglie *et al.*, Science 224:838-843 (1984)); or heat shock promoters, e.g., soybean hsp17.5-E or hsp17.3-B (Gurley *et al.*, Mol. Cell. Biol., 6:559-565 (1986)) may be used. These constructs can be introduced into plant cells using Ti plasmids, Ri plasmids, plant virus vectors, direct DNA transformation, microinjection, electroporation, or other techniques that are well known in the art. For reviews of such techniques see, for example, Weissbach and Weissbach, Methods for Plant Molecular Biology,



Academic Press, NY, Section VIII, pp. 421-463 (1988); and Grierson and Corey, Plant Molecular Biology, 2d Ed., Blackie, London, Ch. 7-9 (1988). Eukaryotic cells are alternative host cells for the expression of mutated *S. cerevisiae* squalene synthase coding sequences. Such host cell lines may include but are not limited to CHO, VERO, BHK, HeLa, COS, MDCK, 293, and WI38.

[0054] Mammalian cell systems that utilize recombinant viruses or viral elements to direct expression may be engineered. For example, when using adenovirus expression vectors, the coding sequence of a variant squalene synthase enzyme may be ligated to an adenovirus transcription/translation control complex, *e.g.*, the late promoter and tripartite leader sequence. This chimeric gene may then be inserted into the adenovirus genome by *in vitro* or *in vivo* recombination. Insertion in a non-essential region of the viral genome (*e.g.*, region E1 or E3) will result in a recombinant virus that is viable and capable of expressing the variant squalene synthase enzyme in infected hosts (Logan and Shenk, PNAS USA 81:3655-3659 (1984)). Alternatively, the vaccinia virus 7.5K promoter may be used. (Mackett *et al.*, PNAS USA, 79:7415-7419 (1982); Mackett *et al.*, J. Virol. 49:857-864 (1984); Panicali, *et al.*, PNAS USA, 79:4927-4931 (1982)). Of particular interest are vectors based on bovine papilloma virus which have the ability to replicate as extrachromosomal elements (Sarver *et al.*, Mol. Cell. Biol. 1:486(1981)). Shortly after entry of this DNA into mouse cells, the plasmid replicates to about 100 to 200 copies per cell. Transcription of the inserted cDNA does not require integration of the plasmid into the host's chromosome, thereby yielding a high level of expression. These vectors can be used for stable expression by including a selectable marker in the plasmid, such as the *neo* gene. Alternatively, the retroviral genome can be modified for use as a vector capable of introducing and directing the expression of the variant squalene synthase enzyme in host cells (Cone and Mulligan, PNAS USA 81:6349-6353 (1984)). High level expression may also be achieved using inducible promoters, including, but not limited to, the metallothionein IIA promoter and heat shock promoters.

[0055] Also within the scope of the invention are host cells including a nucleic acid segment encoding the variant squalene synthase according to the present invention, as described above. These host cells are typically transformed or transfected with the nucleic acid segment; methods for such transformation or transfection are described

above. The term "nucleic acid segment" is used herein to include the following alternatives: (1) a vector including therein the variant squalene synthase; or (2) a chromosome of the host cell including therein the variant squalene synthase. The vector or chromosome can include, as described above, nucleic acid sequences either 5'-, 3'-, or both 5'- and 3'- to the coding sequence of the mutated *S. cerevisiae* squalene synthase, such as, but not limited to, linkers, adaptors, restriction endonuclease cleavage sites, regulatory sequences such as promoters, enhancers, or operators, or coding sequences. The host cells can be prokaryotic cells, such as bacteria, or can be eukaryotic cells, such as yeast cells, plant cells, or animal cells. If the host cells are yeast cells, they are typically *S. cerevisiae*, although other genera of yeast, such as *Pichia* (*Pichia pastoris*) or *Kluyveromyces* (*Kluyveromyces lactis*) can also be employed. If the cells are plant cells, many types of plant cells are suitable host cells; one frequently employed host cell is *Arabidopsis thaliana*. If the cells are animal cells, they can be insect cells or mammalian cells.

[0056] In host cells according to the present invention including therein a nucleic acid segment encoding the variant squalene synthase as described above, the nucleic acid segment can be incorporated into a vector as described above. Alternatively, the nucleic acid segment can be integrated into a chromosome of the host cell. Methods for integrating the nucleic acid segment encoding the variant squalene synthase as described above into the chromosome of a prokaryotic cell (*i.e.*, a bacterium) or into one chromosome of a eukaryotic cell are known in the art. As described above, in this application, the nucleic acid is typically DNA.

[0057] DNA sequences encoding variant squalene synthase can be obtained by several methods. For example, the DNA can be isolated using hybridization procedures that are well known in the art. These include, but are not limited to: (1) hybridization of probes to genomic or cDNA libraries to detect shared nucleotide sequences; (2) antibody screening of expression libraries to detect shared structural features; and (3) synthesis by the polymerase chain reaction (PCR). RNA sequences of the invention can be obtained by methods known in the art (See, for example, Current Protocols in Molecular Biology, Ausubel, *et al.*, Eds. (1989)).

[0058] The development of specific DNA sequences encoding variant squalene synthases of the invention can be obtained by: (1) isolation of a double-stranded DNA sequence from the genomic DNA; (2) chemical manufacture of a DNA sequence to provide the necessary codons for the polypeptide of interest; and (3) in vitro synthesis of a double-stranded DNA sequence by reverse transcription of mRNA isolated from a eukaryotic donor cell. In the latter case, a double-stranded DNA complement of mRNA is eventually formed which is generally referred to as cDNA. Of these three methods for developing specific DNA sequences for use in recombinant procedures, the isolation of genomic DNA is the least common. This is especially true when it is desirable to obtain the microbial expression of eukaryotic polypeptides, such as yeast polypeptides, due to the presence of introns. For obtaining nucleic acid sequences encoding variant squalene synthases according to the present invention, the synthesis of DNA sequences is frequently the method of choice when the entire sequence of amino acid residues of the desired polypeptide product is known. When the entire sequence of amino acid residues of the desired polypeptide is not known, the direct synthesis of DNA sequences is not possible and the method of choice is the formation of cDNA sequences. Among the standard procedures for isolating cDNA sequences of interest is the formation of plasmid-carrying cDNA libraries which are derived from reverse transcription of mRNA which is abundant in donor cells that have a high level of genetic expression. When used in combination with polymerase chain reaction technology, even rare expression products can be clones. In those cases where significant portions of the amino acid sequence of the polypeptide are known, the production of labeled single or double-stranded DNA or RNA probe sequences duplicating a sequence putatively present in the target cDNA may be employed in DNA/DNA hybridization procedures which are carried out on cloned copies of the cDNA which have been denatured into a single-stranded form (Jay *et al.*, Nucleic Acid Research 11:2325 (1983)).

[0059] Nucleotide sequences encompassed by the present invention can also be incorporated into a vector as described above, including, but not limited to, an expression vector, and used to transfect or transform suitable host cells, as is well known in the art. The vectors incorporating the nucleotide sequences that are encompassed by the present invention are also within the scope of the invention. Host cells that are transformed or

transfected with the vector or with polynucleotides or nucleotide sequences of the present invention are also within the scope of the invention. The host cells can be prokaryotic or eukaryotic; if eukaryotic, the host cells can be mammalian cells, insect cells, or yeast cells. If prokaryotic, the host cells are typically bacterial cells.

[0060] Transformation of a host cell with recombinant DNA may be carried out by conventional techniques as are well known to those skilled in the art. Where the host is prokaryotic, such as *Escherichia coli*, competent cells which are capable of DNA uptake can be prepared from cells harvested after exponential growth phase and subsequently treated by the  $\text{CaCl}_2$  method by procedures well known in the art. Alternatively,  $\text{MgCl}_2$  or  $\text{RbCl}$  can be used. Transformation can also be performed after forming a protoplast of the host cell or by electroporation.

[0061] When the host is a eukaryote, such methods of transfection of DNA as calcium phosphate co-precipitates, conventional mechanical procedures such as microinjection, electroporation, insertion of a plasmid encased in liposomes, or virus vectors may be used.

[0062] A variety of host-expression vector systems may be utilized to express the nucleic acid sequence encoding the variant squalene synthase enzymes of the present invention. These include but are not limited to microorganisms such as bacteria transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing a variant squalene synthase enzyme coding sequence; yeast transformed with recombinant yeast expression vectors containing the variant squalene synthase enzyme coding sequence; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing a variant squalene synthase enzyme coding sequence; insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus) containing a variant squalene synthase enzyme coding sequence; or animal cell systems infected with recombinant virus expression vectors (e.g., retroviruses, adenovirus, vaccinia virus) containing a variant squalene synthase enzyme coding sequence, or transformed animal cell systems engineered for stable expression. In such cases where glycosylation may be important,

expression systems that provide for translational and post-translational modifications may be used; *e.g.*, mammalian, insect, yeast or plant expression systems.

[0063] Depending on the host/vector system utilized, any of a number of suitable transcription and translation elements, for example, constitutive and inducible promoters, transcription enhancer elements, transcription terminators, may be used in the expression vector (Bitter *et al.*, *Methods in Enzymology*, 153:516-544 (1987)). For example, when cloning in bacterial systems, inducible promoters such as pL of bacteriophage  $\lambda$ , plac, ptrp, ptac (ptrp-lac hybrid promoter) and the like may be used. When cloning in mammalian cell systems, promoters derived from the genome of mammalian cells (*e.g.*, metallothionein promoter) or from mammalian viruses (*e.g.*, the retrovirus long terminal repeat; the adenovirus late promoter; the vaccinia virus 7.5K promoter) may be used. Promoters produced by recombinant DNA or synthetic techniques may also be used to provide for transcription of the inserted variant squalene synthase enzyme coding sequence.

[0064] In bacterial systems a number of expression vectors may be advantageously selected depending upon the use intended for the variant squalene synthase enzyme expressed, and whether it is desired to isolate the enzyme and in what state of purity. For example, when large quantities are to be produced, vectors which direct the expression of high levels of fusion protein products that are readily purified may be desirable. Those which are engineered to contain a cleavage site to aid in recovering the protein are preferred. Such vectors include, but are not limited to, the *Escherichia coli* expression vector pUR278 (Ruther *et al.*, *EMBO J.*, 2:1791 (1983)), in which the variant squalene synthase enzyme coding sequence may be ligated into the vector in frame with the lac Z coding region so that a hybrid variant squalene synthase enzyme-lac Z protein is produced as well as pIN vectors (Inouye and Inouye, *Nucleic Acids Res.* 13:3101-3109 (1985); Van Heeke and Schuster, *J. Biol. Chem.* 264:5503-5509 (1989)).

[0065] In yeast, a number of vectors containing constitutive or inducible promoters may be used. For a review, see *Current Protocols in Molecular Biology*, Vol. 2, Ed. Ausubel, *et al.*, Greene Publish. Assoc. & Wiley Interscience, Ch. 13 (1988); Grant *et al.*, *Expression and Secretion Vectors for Yeast*, in *Methods in Enzymology*, Eds. Wu

& Grossman, 31987, Acad. Press, N.Y., Vol. 153, pp.516-544 (1987); Glover, DNA Cloning, Vol. II, IRL Press, Wash., D.C., Ch. 3 (1986); Bitter, Heterologous Gene Expression in Yeast, Methods in Enzymology, Eds. Berger & Kimmel, Acad. Press, N.Y., Vol. 152, pp. 673-684 (1987); and The Molecular Biology of the Yeast *Saccharomyces*, Eds. Strathern et al., Cold Spring Harbor Press, Vols. I and II (1982). A constitutive yeast promoter such as ADH1 or LEU2 or an inducible promoter such as GAL4 may be used (Cloning in Yeast, Ch. 3, R. Rothstein In: DNA Cloning Vol. II, A Practical Approach, Ed. DM Glover, IRL Press, Wash., D.C. (1986)). Alternatively, vectors may be used which promote integration of foreign DNA sequences into the yeast chromosome.

[0066] In cases where plant expression vectors are used, the expression of a variant squalene synthase enzyme coding sequence may be driven by any of a number of promoters. For example, viral promoters such as the 35S RNA and 19S RNA promoters of CaMV (Brisson *et al.*, Nature, 310:511-514 (1984)), or the coat protein promoter to TMV (Takamatsu *et al.*, EMBO J., 6:307-311 (1987)) may be used; alternatively, plant promoters such as the small subunit of RUBISCO (Coruzzi *et al.*, EMBO J. 3:1671-1680 (1984); Broglie *et al.*, Science 224:838-843 (1984)); or heat shock promoters, e.g., soybean hsp17.5-E or hsp17.3-B (Gurley *et al.*, Mol. Cell. Biol., 6:559-565 (1986)) may be used. These constructs can be introduced into plant cells using Ti plasmids, Ri plasmids, plant virus vectors, direct DNA transformation, microinjection, electroporation, or other techniques that are well known in the art. (Weissbach & Weissbach, Methods for Plant Molecular Biology, Academic Press, NY, Section VIII, pp. 421-463 (1988); Grierson and Corey, Plant Molecular Biology, 2d Ed., Blackie, London, Ch. 7-9 (1988).

[0067] An alternative expression system that can be used to express a variant squalene synthase enzyme of the present invention is an insect system. In one such system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes. The virus grows in *Spodoptera frugiperda* cells. The variant squalene synthase enzyme coding sequence may be cloned into non-essential regions (in *Spodoptera frugiperda*, for example, the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter). Successful insertion of the variant squalene synthase enzyme coding sequence will result in

inactivation of the polyhedrin gene and production of non-occluded recombinant virus (*i.e.*, virus lacking the proteinaceous coat coded for by the polyhedrin gene). These recombinant viruses are then used to infect cells in which the inserted gene is expressed. (Smith *et al.*, J. Biol. 46:584 (1983); U.S. Pat. No. 4,215,051).

[0068] Eukaryotic systems, and preferably mammalian expression systems, allow for proper post-translational modifications of expressed eukaryotic proteins to occur. Therefore, eukaryotic cells, such as mammalian cells that possess the cellular machinery for proper processing of the primary transcript, glycosylation, phosphorylation, and, advantageously secretion of the gene product, are the preferred host cells for the expression of a variant squalene synthase enzyme. Such host cell lines may include, for example CHO, VERO, BHK, HeLa, COS, MDCK, 293, and WI38. Other eukaryotic host cells are also suitable host cells for the expression of a variant squalene synthase enzyme or of nucleic acids according to the present invention, including, for example, microalgal cells.

[0069] Mammalian cell systems that utilize recombinant viruses or viral elements to direct expression may be engineered for use in the present invention. For example, when using adenovirus expression vectors, the coding sequence of a variant squalene synthase enzyme may be ligated to an adenovirus transcription/translation control complex, *e.g.*, the late promoter and tripartite leader sequence. This chimeric gene may then be inserted into the adenovirus genome by *in vitro* or *in vivo* recombination. Insertion in a non-essential region of the viral genome (*e.g.*, region E1 or E3) will result in a recombinant virus that is viable and capable of expressing the variant squalene synthase enzyme in infected hosts (Logan and Shenk, PNAS USA 81:3655-3659 (1984)). Alternatively, the vaccinia virus 7.5K promoter may be used (Mackett *et al.*, PNAS USA, 79:7415-7419 (1982); Mackett *et al.*, J. Virol. 49:857-864, 1984; Panicali *et al.*, (PNAS USA, 79:4927-4931 (1982)). Of particular interest are vectors based on bovine papilloma virus which have the ability to replicate as extrachromosomal elements (Sarver *et al.*, Mol. Cell. Biol. 1:486 (1981)). Shortly after entry of this DNA into mouse cells, the plasmid replicates to about 100 to 200 copies per cell. Transcription of the inserted cDNA does not require integration of the plasmid into the host's chromosome, thereby yielding a high level of expression. These vectors can be used for stable expression by

including a selectable marker in the plasmid, such as the *neo* gene. Alternatively, the retroviral genome can be modified for use as a vector capable of introducing and directing the expression of the variant squalene synthase enzyme in host cells (Cone and Mulligan, PNAS USA 81:6349-6353 (1984)). High level expression may also be achieved using inducible promoters, including, but not limited to, the metallothionein IIA promoter and heat shock promoters.

[0070] For long-term, high-yield production of recombinant proteins, stable expression is preferred. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with a cDNA controlled by appropriate expression control elements (e.g., promoter, enhancer, sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines. For example, following the introduction of foreign DNA, engineered cells may be allowed to grow for 1-2 days in enriched media, and then are switched to a selective medium. A number of selection systems may be used, including but not limited, to the herpes simplex virus thymidine kinase (Wigler *et al.*, Cell 11:223 (1977)), hypoxanthine-guanine phosphoribosyltransferase (Szybalska and Szybalski, PNAS USA, 48:2026 (1962)), and adenine phosphoribosyltransferase (Lowy *et al.*, Cell, 22:817 (1980)) genes, which can be employed in *tk<sup>-</sup>*, *hgprt<sup>-</sup>* or *aprt<sup>-</sup>* cells, respectively. Additionally, antimetabolite resistance-conferring genes can be used as the basis of selection; for example, the genes for *dhfr*, which confer resistance to methotrexate (Wigler *et al.*, Natl. Acad. Sci. USA, 77:3567 (1980); O'Hare *et al.*, PNAS USA, 78:1527 (1981)); *gpt*, which confers resistance to mycophenolic acid (Mulligan and Berg, PNAS USA, 78:2072 (1981)); *neo*, which confers resistance to the aminoglycoside G418 (Colberre-Garapin *et al.*, J. Mol. Biol., 150:1 (1981)); and *hygro*, which confers resistance to hygromycin (Santerre *et al.*, Gene, 30:147 (1984)). Recently, additional selectable genes have been described, namely *trpB*, which allows cells to utilize indole in place of tryptophan; *hisD*, which allows cells to utilize histinol in place of histidine (Hartman and Mulligan, PNAS USA, 85:804 (1988)); and ODC (ornithine decarboxylase) which confers resistance to the ornithine decarboxylase inhibitor, 2-



(difluoromethyl)-DL-ornithine, DFMO (McConlogue, In: Current Communications in Molecular Biology, Cold Spring Harbor Laboratory ed., (1987)).

[0071] Another aspect of the present invention is a variant squalene synthase enzyme encoded by a nucleic acid segment of the present invention as described above. These squalene synthase enzymes can be, for example, enzymes derived from *S. cerevisiae* or other species of the genus *Saccharomyces*; they can also include other squalene synthases derived from any organism that has a gene that catalyzes the conversion of farnesyl pyrophosphate into squalene.

[0072] When the variant squalene synthase enzymes are *S. cerevisiae* squalene synthase enzymes, they include, for example, the following:

- (1) a variant *S. cerevisiae* squalene synthase enzyme as shown as Mutant 4 in Table 2;
- (2) a variant *S. cerevisiae* squalene synthase enzyme as shown as Mutant 10 in Table 2;
- (3) a variant *S. cerevisiae* squalene synthase enzyme as shown as Mutant 14 in Table 2;
- (4) a variant *S. cerevisiae* squalene synthase enzyme as shown as Mutant 19 in Table 2;
- (5) a variant *S. cerevisiae* squalene synthase enzyme as shown as Mutant 22 in Table 2;
- (6) a variant *S. cerevisiae* squalene synthase enzyme as shown as Mutant 23 in Table 2;
- (7) a variant *S. cerevisiae* squalene synthase enzyme as shown as Mutant 24 in Table 2;
- (8) a variant *S. cerevisiae* squalene synthase enzyme as shown as Mutant 25 in Table 2;
- (9) a variant *S. cerevisiae* squalene synthase enzyme as shown as Mutant 69 in Table 2;
- (10) a variant *S. cerevisiae* squalene synthase enzyme in which the wild-type *S. cerevisiae* squalene synthase enzyme is mutated with the same amino acid changes as in any of (1) through (9) above;

(11) a variant *S. cerevisiae* squalene synthase enzyme containing any of the amino acid changes as in any of (1) through (10), above; and

(12) a variant *S. cerevisiae* squalene synthase enzyme in which the squalene synthase enzyme differs from the variant squalene synthase enzyme of any of (1) through (11) above by one to three conservative amino acid substitutions, wherein a conservative amino acid substitution is defined as one of the following substitutions: A→G or S; R→K; N→Q or H; D→E; C→S; Q→N; G→D; G→A or P; H→N or Q; I→L or V; L→I or V; K→R or Q or E; M→L or Y or I; F→M or L or Y; S→T; T→S; W→Y; Y→W or F; and V→I or L.

[0073] Furthermore, with respect to these alternatives, typically, the variant squalene synthase enzyme has a reduced  $V_{\max}$  for squalene synthesis. Alternatively, the variant squalene synthase enzyme has an increased  $K_m$  for its FPP substrate, in which case the enzyme is less active at a given intracellular concentration of FPP than the wild-type enzyme. Typically, the variant squalene synthase enzyme, when expressed *in vivo* in a suitable eukaryotic microbial host, as described above, produces squalene at a rate of less than 75% of the wild-type enzyme. Preferably, the variant squalene synthase enzyme, when expressed *in vivo* in a suitable eukaryotic microbial host, as described above, produces squalene at a rate of less than 50% of the wild-type enzyme. More preferably, the variant squalene synthase enzyme, when expressed *in vivo* in a suitable eukaryotic microbial host, produces squalene at a rate of less than 25% of the wild-type enzyme.

[0074] With respect to (12) of these alternatives, preferably, the variant *S. cerevisiae* squalene synthase enzyme differs from the variant *S. cerevisiae* squalene synthase enzyme of any of (1) through (10) above by one to two conservative amino acid substitutions. More preferably, the variant *S. cerevisiae* squalene synthase enzyme differs from the variant *S. cerevisiae* squalene synthase enzyme of any of (1) through (10) above by one conservative amino acid substitution.

[0075] Another aspect of the present invention is a host cell containing and/or expressing a variant squalene synthase enzyme of the present invention as described above. The host cell, in this alternative, includes at least one copy of a nucleic acid sequence encoding a variant squalene synthase enzyme. The at least one copy of the

nucleic acid sequence encoding a variant squalene synthase enzyme can be present in the chromosome of a prokaryotic (bacterial) cell or in one chromosome of a eukaryotic cell. Alternatively, the at least one copy of the nucleic acid sequence encoding a variant squalene synthase enzyme can be present in a vector or plasmid that is present in the cell. The host cell, as described above, can be a prokaryotic or eukaryotic cell. If a prokaryotic cell, it can be a bacterial cell. If a eukaryotic cell, it can be a yeast cell, a plant cell, or an animal cell. Suitable host cells are described above.

[0076] Another aspect of the present invention is a method of isolating a mutated *ERG9* gene. The mutated *ERG9* gene is typically a *S. cerevisiae* gene, but can be a homologous gene from another species as described above.

[0077] In general, a method of isolating a mutated *ERG9* gene according to the present invention comprises the steps of:

- (1) isolating a wild-type *ERG9* gene to produce an isolated wild-type *ERG9* gene;
- (2) subjecting the isolated wild-type *ERG9* gene to mutagenesis to generate a pool of *ERG9* mutants;
- (3) transforming mutants from the pool of *ERG9* mutants generated in step (b) into a strain of a eukaryotic microbial host that contains a plasmid expressing a terpene synthase gene that produces a detectable and measurable terpene product, the strain of the eukaryotic microbial host being transformed in such a manner that replacement of the preexisting *erg9* allele with an *erg9* mutation allows the strain to grow in a sterol-free medium; and
- (4) isolating a transformant from step (c) that produces a level of terpene product at least equivalent to the level of terpene product produced by a strain of the eukaryotic microbial host expressing the terpene synthase gene that requires a sterol in the medium for growth.

[0078] In the present invention, the step of isolating a wild-type *ERG9* gene to produce an isolated wild-type *ERG9* gene is typically performed by amplifying a wild-type *ERG9* gene by using a nucleic acid amplification process to produce an amplified wild-type *ERG9* gene; however, other isolation methods are known in the art and are contemplated by this invention, it is not necessary to use PCR or another nucleic acid

amplification method. When a nucleic acid amplification process is used, the nucleic acid amplification method is typically PCR. However, other nucleic acid amplification processes can be used that are well known in the art. In this method, the *ERG9* gene can be a fungal *ERG9* gene, such as a *Saccharomyces cerevisiae* *ERG9* gene or an *ERG9* gene of another *Saccharomyces* species; alternatively, the *ERG9* gene can be any homologous *ERG9* gene as described above. In this method, the mutagenesis is typically performed using error-prone PCR, although other mutagenesis methods are well known in the art and can alternatively be used. Such mutagenesis methods include, for example, ultraviolet (UV) radiation, ethyl methanesulfonate (EMS), nitrosoguanidine, and other mutagens.

[0079] When error-prone PCR is used, the error-prone PCR is typically performed using one or more DNA polymerase enzymes that have higher misinsertion and misextension rates than wild-type polymerase enzymes.

[0080] In the present invention, the terpene synthase gene that produces a detectable and measurable terpene product can be, for example, the *Hyoscyamus muticus* premnaspirodiene synthase (HPS) gene. Preferably, the terpene synthase gene that produces a detectable and measurable terpene product is one that produces a product detectable and measurable by gas chromatography, although other detection methods can be used.

[0081] In the present invention, a suitable strain of *S. cerevisiae* that contains a plasmid expressing a terpene synthase gene that produces a detectable and measurable terpene product is ALX7-95 (*his3, trp1, erg9::HIS3, HMGcat/TRP1::rDNA, dpp1*), a leucine prototroph of strain CALI-5 (US patents 6,531,303 and 6,689,593), containing a plasmid expressing the *Hyoscyamus muticus* premnaspirodiene synthase (HPS) gene. Transformants of this strain require histidine for growth; before transformation, this strain requires supplementation with a sterol.

[0082] Although the invention can be used with isolates of mutations in *S. cerevisiae*, analogous methods can be used with other organisms, as described above.

[0083] Another aspect of the present invention is a method of isolating a variant squalene synthase enzyme. The variant squalene synthase enzyme to be isolated by these methods is as described above.

[0084] In general, this method comprises the steps of:

- (1) culturing a host cell that expresses a variant squalene synthase enzyme or that contains a variant squalene synthase enzyme; and
- (2) isolating the variant squalene synthase enzyme from the host cell.

[0085] Typically, this method further comprises the step of purifying the isolated variant squalene synthase enzyme. Variant squalene synthase enzymes according to the present invention can be purified by conventional protein purification techniques, including, for example, techniques such as precipitation with salts such as ammonium sulfate, ion exchange chromatography, gel filtration, affinity chromatography, electrophoresis, isoelectric focusing, isotachopheresis, chromatofocusing, and other techniques are well known in the art and are described in R.K. Scopes, "Protein Purification: Principles and Practice" (3<sup>rd</sup> ed., Springer-Verlag, New York (1994)).

[0086] Yet another aspect of the present invention is a method of producing an isoprenoid using a mutated *ERG9* gene, in which the defective *ERG9* gene encodes a variant squalene synthase enzyme.

[0087] In one aspect of the invention, a host cell that includes a mutated *ERG9* gene encoding a variant squalene synthase enzyme further includes at least one isoprenoid synthase gene, so that the farnesyl pyrophosphate produced in the host cell, which is available in greater concentrations for isoprenoid biosynthesis can be converted to an isoprenoid by the isoprenoid synthase encoded by the isoprenoid synthase gene. For example, and not by way of limitation, the isoprenoid synthase gene included in the host cell can be a chimeric isoprenoid synthase gene such as those described in U.S. Patent Application Publication No. 2008/0178354. These chimeric isoprenoid synthase genes include derivatives of the *Hyoscyamus muticus* vetispiradiene synthase gene and/or the *Nicotiana tabacum* 5-epi-aristolochene synthase gene. Alternatively, the isoprenoid synthase gene included in the host cell can be a citrus valencene synthase gene as described in U.S. Patent Application Publication No. 2006/0218661. As yet another alternative, the isoprenoid synthase gene included in the host cell can be a *H. muticus* premnaspirodiene synthase gene, such as those described in Back and Chappell, J. Biol. Chem. 270: 7375-7381 (1995); Back and Chappell, PNAS USA 93: 6841-6845 (1996); and Greenhagen *et al.*, PNAS USA 103: 9826-9831 (2006). As another alternative, the

isoprenoid synthase gene included in the host cell can be an isoprenoid synthase gene such as those described in U.S. Patent Application Publication No. 2005/0210549. These isoprenoid synthase genes include 5-epi-aristolochene synthase from *Capsicum annuum*, (E)- $\beta$ -farnesene synthase from *Mentha piperita*,  $\delta$ -selenene synthase and  $\gamma$ -humulene synthase from *Abies grandis*, 6-cadinene synthase from *Gossypium arboreum*, E- $\alpha$ -bisabolene synthase from *Abies grandis*, germacrene C synthase from *Lycopersicon esculentum*, epi-cedrol synthase and amorpho-4,11-diene synthase from *Artemisia annua*, and germacrene A synthases from *Lactuca sativa*, *Cichorium intybus* and *Solidago canadensis*. Other suitable isoprenoid synthase genes are known in the art. In addition, mutants and protein engineered variants of these enzymes can be used. Methods for engineering variants of terpene synthases are known in the art; such methods can, for example, involve recombining domains from two or more terpene synthases to generate a chimeric terpene synthase. Such methods are described, for example, in U.S. Patent Application Publication No. 2006/0218661 and in U.S. Patent Application Publication No. 2008/0178354.

[0088] Accordingly, this aspect of the invention comprises the steps of:

- (1) providing a host cell including a mutated *ERG9* gene according to the present invention and at least one isoprenoid synthase gene;
- (2) allowing the host cell to produce farnesyl pyrophosphate and to synthesize the isoprenoid from the farnesyl pyrophosphate; and
- (3) isolating the isoprenoid synthesized by the host cell.

[0089] Methods for isolating the isoprenoid synthesized by the host cell are well known in the art. For example, when the isoprenoid is premnaspirodiene, the premnaspirodiene can be isolated by (i) sequestering the premnaspirodiene by binding it to a hydrophobic resin; (ii) and isolating the premnaspirodiene from the hydrophobic resin. Preferably, the hydrophobic resin is Amberlite® XAD-16 hydrophobic resin. Other hydrophobic resins within the scope of the present invention will be known to one of reasonable skill in the art. Typically, premnaspirodiene is isolated from the hydrophobic resin by methanol extraction. Other methods of isolating premnaspirodiene that are within the scope of the present invention will be known to one of reasonable skill

in the art. Other isoprenoids can be isolated by similar methods well known in the art, making use of the fact that isoprenoids are hydrophobic and bind to hydrophobic resins.

[0090] In an alternative for isolation of premnaspirodiene, a two-phase system can be used with a non-polar solvent, substantially immiscible with an aqueous phase, added to the fermentation broth and the premnaspirodiene removed from the non-polar phase by distillation. A preferred non-polar solvent is an oil. A particularly preferred oil is a vegetable oil such as soybean oil. Alternative non-polar solvents include, but are not limited to, high molecular weight aliphatic hydrocarbons such as, but not limited to, dodecane, tridecane, tetradecane, pentadecane, and hexadecane; either straight-chain or branched-chain isomers can be used; these high-molecular weight aliphatic hydrocarbons are optionally substituted with one or more hydroxy or halogen substituents as long as the substituted hydrocarbon remains substantially immiscible with the aqueous phase.

[0091] In this method, where commercial production of the isoprenoid is desired, a variety of fermentation methodologies may be applied. For example, large scale production may be effected by either batch or continuous fermentation. A classical batch fermentation is a closed system where the composition of the medium is set at the beginning of the fermentation and not subject to artificial alterations during the fermentation. Thus, at the beginning of the fermentation the medium is inoculated with the desired microorganism or microorganisms and fermentation is permitted to occur without further addition of nutrients. Typically, the concentration of the carbon source in a batch fermentation is limited, and factors such as pH and oxygen concentration are controlled. In batch systems the metabolite and biomass compositions of the system change constantly up to the time the fermentation is stopped. Within batch cultures cells typically modulate through a static lag phase to a high growth log phase and finally to a stationary phase where growth rate is diminished or halted. If untreated, cells in the stationary phase will eventually die.

[0092] A variation on the standard batch system is the Fed-Batch system. Fed-Batch fermentation processes are also suitable for use in the present invention and comprise a typical batch system with the exception that nutrients are added as the fermentation progresses. Fed-Batch systems are useful when catabolite repression is apt to inhibit the metabolism of the cells and where it is desirable to have limited amounts of

substrate in the medium. Also, the ability to feed nutrients will often result in higher cell densities in Fed-Batch fermentation processes compared to Batch fermentation processes. Factors such as pH, dissolved oxygen, nutrient concentrations, and the partial pressure of waste gases such as CO<sub>2</sub> are generally measured and controlled in Fed-Batch fermentations. Batch and Fed-Batch fermentations are common and well known in the art and examples may be found in Brock, *Biotechnology: A Textbook of Industrial Microbiology*, 2nd ed.; Sinauer Associates: Sunderland, Mass. (1989); or Deshpande, *Appl. Biochem. Biotechnol.* 36:227 (1992).

[0093] Commercial production of the isoprenoid may also be accomplished with continuous fermentation. Continuous fermentation is an open system where a defined fermentation medium is added continuously to a bioreactor and an equal amount of conditioned medium is removed simultaneously for processing. This system generally maintains the cultures at a constant high density where cells are primarily in their log phase of growth. Continuous fermentation allows for modulation of any number of factors that affect cell growth or end product concentration. For example, one method will maintain a limiting nutrient such as the carbon source or nitrogen level at a fixed rate and allow all other parameters to moderate. In other systems a number of factors affecting growth can be altered continuously while the cell concentration, measured by the medium turbidity, is kept constant. Continuous systems strive to maintain steady state growth conditions and thus the cell loss due to the medium removal must be balanced against the cell growth rate in the fermentation. Methods of modulating nutrients and growth factors for continuous fermentation processes as well as techniques for maximizing the rate of product formation are well known in the art of industrial microbiology and a variety of methods are detailed by Brock, *supra*.

[0094] Microorganism host cells useful in the present invention for the production of the isoprenoid may include, but are not limited to, bacteria, such as the enteric bacteria (*Escherichia* and *Salmonella* for example) as well as *Bacillus*, *Acinetobacter*, *Streptomyces*, *Methylobacter*, *Rhodococcus* and *Pseudomonas*; Cyanobacteria, such as *Rhodobacter* and *Synechocystis*; yeasts, such as *Saccharomyces*, *Zygosaccharomyces*, *Kluyveromyces*, *Candida*, *Hansenula*, *Debaryomyces*, *Mucor*, *Pichia*, *Yarrowia*, and *Torulopsis*; and filamentous fungi such as *Aspergillus* and



*Arthrobotrys*, and algae for example. Preferably, the host cell is a eukaryotic cell. More preferably, the host cell is a yeast cell, which is a eukaryotic microorganism host cell. Most preferably, the host cell is a *Saccharomyces cerevisiae* cell.

[0095] Microbial expression systems and expression vectors containing regulatory sequences that direct high level expression of foreign proteins are well known to those skilled in the art. These expression systems and expression vectors are known both for prokaryotic organisms such as bacteria and for eukaryotic organisms such as yeast. Similarly, vectors or cassettes useful for the transformation of suitable microbial host cells are well known in the art. These vectors and cassettes are known both for prokaryotic organisms such as bacteria and for eukaryotic organisms such as yeast. Typically, the vector or cassette contains sequences directing expression of the relevant gene, a selectable marker, and sequences allowing autonomous replication or chromosomal integration. Suitable vectors comprise a region 5' of the gene which harbors transcriptional initiation controls and a region 3' of the DNA fragment which controls transcriptional termination.

[0096] Initiation control regions or promoters, which are useful to drive expression of the relevant genes in the desired host cell are numerous and familiar to those skilled in the art. Termination control regions may also be derived from various genes native to the preferred hosts.

[0097] Expression of cloned heterologous genes in yeast cells, particularly cells of *S. cerevisiae*, is described in the following references: Emr, Meth. Enzymol. 185: 231-233 (1991), is a general overview of expression in yeast, including the possibility of exploiting protein secretion and modification in yeast and achieving stability of expressed proteins. Rose and Broach, Meth. Enzymol. 185: 234-279 (1991), describes the use of 2- $\mu$ m circle-based vectors for transfection of genes into yeast and for expression of heterologous genes in yeast, including standard 2- $\mu$ m circle-based vectors, vectors for high copy propagation, vectors for expression of cloned genes in yeast, and vectors for specialized applications. Stearns *et al.*, Meth. Enzymol. 185: 280-297 (1991), describes the use of yeast vector systems and components, the use of homologous recombination to integrate plasmids into the yeast host genome, and the use of centromere plasmids. Mylin *et al.*, Meth. Enzymol. 185: 297-308 (1991), describes the use of galactose-

inducible promoters to provide high levels of production of cloned proteins in yeast. Price *et al.*, Meth. Enzymol. 185: 308-318 (1991), describes the use of the glucose-repressible *ADH2* promoter to provide controllable, high level expression of cloned proteins in yeast. Etcheverry, Meth. Enzymol. 185: 319-329 (1991), describes the use of the yeast *CUP1* promoter to drive controllable expression of cloned genes in yeast. Kingsman *et al.*, Meth. Enzymol. 185: 329-341 (1991), describes the use of the yeast *PGK* promoter to drive controllable expression of cloned genes in yeast. Rosenberg *et al.*, Meth. Enzymol. 185: 341-350 (1991), describes the use of expression cassette plasmids utilizing the strong *GAPDH-491* promoter for high levels of heterologous protein production in yeast. Sledziewski *et al.*, Meth. Enzymol. 185: 351-366 (1991), describes the construction of temperature-regulated variants of two strong yeast promoters, *TPII* and *ADH2*, and the use of these promoters for regulation of expression and thus regulation of the extent of glycosylation of proteins secreted by yeast. Donahue and Cigan, Meth. Enzymol. 185: 366-372 (1991), describes the significance of codon usage variations between yeast and higher eukaryotes and the selection of efficient leader sequences. Jones, Meth. Enzymol. 185: 372-386 (1991), describes the elimination of vacuolar protease activity in yeast to maximize the yield of protein production from cloned genes. Wilkinson, Meth. Enzymol. 185: 387-397 (1991), describes methods for preventing ubiquitin-dependent protein degradation in yeast, again to maximize the yield of protein production from cloned genes. Kendall *et al.*, Meth. Enzymol. 185: 398-407 (1991), describes the cotranslational processing events that occur in yeast at the amino-termini of nascent polypeptide genes and their effects on heterologous gene expression and protein stability. Brake, Meth. Enzymol. 185: 408-421 (1991), describes expression systems based on the yeast  $\alpha$ -factor leader. Hitzeman *et al.*, Meth. Enzymol. 185: 421-440 (1991), describes the use of both heterologous and homologous signal sequences for the production and secretion of heterologous gene products in yeast. Chisholm *et al.*, "Meth. Enzymol. 185: 471-482 (1991), describes the use of an enhanced secretion phenotype occurring among drug-resistant yeast mutants to maximize secretion of cloned proteins in yeast.

[0098] General molecular biological techniques of gene cloning, site-directed mutagenesis, and fusion protein construction can be used to provide nucleic acid

segments that include therein the isoprenoid synthase gene. Typically, the nucleic acid segments are DNA nucleic acid segments. Typically, as described above, the isoprenoid synthase gene is operatively linked to at least one nucleic acid expression control element, such as, but not limited to, a promoter, an enhancer, or a site capable of binding a repressor or activator. Such nucleic acid expression control elements are well known in the art. Typically, as described above, the isoprenoid synthase gene is included in a vector and, as such, is again operatively linked to at least one nucleic acid expression control element. Site-directed mutagenesis can be used, for example, to provide optimum codon selection for expression in *S. cerevisiae*, as described above. The isoprenoid synthase gene can, in one alternative, be expressed in the form of a nucleic acid segment encoding a fusion protein, such as a purification tag or other detectable protein domain.

[0099] In another alternative method for producing an isoprenoid, the method, in general, comprises the steps of:

- (1) providing a host cell including a mutated *ERG9* gene according to the present invention;
- (2) allowing the host cell to produce farnesyl pyrophosphate
- (3) isolating farnesyl pyrophosphate from the host cell;
- (4) reacting the farnesyl pyrophosphate *in vitro* with one or more isoprenoid synthases to synthesize the isoprenoid; and
- (5) isolating the isoprenoid.

[0100] As described above, a number of isoprenoid synthases are available for *in vitro* use. These isoprenoid synthases have been either cloned or isolated from plants. The step of isolating the isoprenoid is performed as described above.

[0101] In both of these synthesis methods, additional reactions can be performed on the isolated isoprenoid to transform the isolated isoprenoid into another isoprenoid or related compound. These additional reactions can be reactions such as oxidation, hydroxylation, alkylation, halogenation, or other reactions well known in the art. In particular, reactions such as hydroxylation or oxidation can be carried out by cytochrome P450 enzymes. These reactions, and methods of carrying them out by chemical or enzymatic means, are well known in the art and need not be described further here.

[0102] The present invention describes improved strains of *Saccharomyces cerevisiae* that have a defective squalene synthase enzyme. These strains have the ability to produce enough squalene so that they do not need to be supplemented with sterols such as ergosterol for growth. However, because these strains produce less squalene than do wild-type strains, they have more farnesyl pyrophosphate available for eventual isoprenoid synthesis, because farnesyl pyrophosphate is a branch point for the steroid synthesis and isoprenoid synthesis pathways. Therefore, these strains, as well as the nucleic acid segments encoding the defective squalene synthase enzyme and the defective squalene synthase enzymes themselves, are useful for the improved production of isoprenoid products because the strains of *S. cerevisiae* do not need to be supplemented with sterols for growth and can produce high levels of farnesyl pyrophosphate without such supplementation.

[0103] Because the eventual isoprenoid products have commercial value as antibiotics, pest control agents, fragrances, flavors, and anti-cancer agents, the nucleic acid segments, eukaryotic microbial host cell strains, including *S. cerevisiae* strains, vectors and host cells incorporating the nucleic acid segments, and the variant squalene synthase enzymes have industrial utility.

[0104] The invention is illustrated by the following examples. These examples are for illustrative purposes only, and are not intended to limit the invention.

## EXAMPLES

### Example 1

#### Generation of Mutant *ERG9* Genes

[0105] Chromosomal DNA isolated from *Saccharomyces cerevisiae* strain ATCC28383 was used as the DNA template for PCR amplification of the wild type *ERG9* gene. The primers used for the amplification were the upstream primer 7-162.1 5'-CCATCTTCAACAACAATACCG-3' (SEQ ID NO: 1) (underlined nucleotides at the 5' end of the *ERG9* sequence in Figure 2) and the downstream primer 7-162.2 5'-GTACTTAGTTATTGTTCCG-3' (SEQ ID NO: 2) (underlined nucleotides at the 3' end of *ERG9*). Using Taq polymerase (New England Biolabs), amplification conditions were

94°C for 30 seconds, 45°C for 30 seconds, 72°C for 2 minutes for a total of 30 cycles. The resulting *ERG9* PCR product was sequenced and verified to be identical to the published sequence for *ERG9*. This DNA was used as the template for performing error prone PCR using the GeneMorph kit from Stratagene and the same primers described above. The error-prone PCR reaction was run using two different DNA concentrations (~500 ng and ~50 ng) to generate a range of mutations per gene. This generated a pool of *ERG9* mutant genes.

### Example 2

#### Isolation of *erg9<sup>def</sup>* Mutants

[0106] To isolate *ERG9* mutants that make sufficient ergosterol to restore growth, the PCR product from the mutagenic PCR reaction was transformed into ALX7-95 (*his3, trp1, erg9::HIS3, HMGcat/TRP1::rDNA, dpp1*), a leucine prototroph of strain CALI-5 (US Patent Nos. 6,531,303 and 6,689,593), containing a plasmid expressing the *Hyoscyamus muticus* premnaspirodiene synthase (HPS) gene. The HPS gene was cloned into the yeast shuttle expression vector YE<sub>p</sub>-GW-URA-NheI/BamHI to give YE<sub>p</sub>-HPS-ura. This vector contained the ADH1 promoter for initiating transcription of the HPS gene. In addition, it contained the ADH1 terminator downstream of the HPS gene. This vector was maintained in *S. cerevisiae* by selecting media lacking uracil and it was maintained in *E. coli* by selecting for resistance to ampicillin.

[0107] Transformation procedure of this strain with the *ERG9* mutant pool used the lithium acetate transformation kit from Sigma. Transformants were selected for growth on YPD medium (10 g/L yeast extract, 20 g/L peptone, 20 g/L glucose) without ergosterol. Since the parent strain requires ergosterol for growth, transformants that grew on YPD replaced the *erg9::HIS3* replacement mutation with a copy of *ERG9* that made sufficient amount of ergosterol for growth. This was verified by the fact that transformants had obtained a requirement for histidine for growth.

[0108] To screen for premnaspirodiene production in strains transformed with *ERG9* mutant genes, a high-throughput screening procedure using microvial cultures was employed. Transformant yeast colonies were inoculated into individual wells of 96-well microtiter plates filled with 200 µL of SD (0.67 Bacto yeast nitrogen base without amino

acids, 2% glucose, 0.14% yeast synthetic drop-out medium without uracil). The plate was grown for two days at 28°C. After growth to saturation, ten  $\mu\text{L}$  from each well was used to inoculate in duplicate two ml glass vials containing 250  $\mu\text{L}$  of medium suitable for growth and premnaspirodiene production. The vials were sealed with serum-stoppered caps and then incubated with shaking for 3 days. The products were extracted first by introducing 250  $\mu\text{L}$  of acetone through the serum stopper and vortexing, followed by addition of 500  $\mu\text{L}$  of *n*-hexane and vortexing. After phase separation, the vials were placed on the sample tray of a gas chromatography autosampler, which removed one microliter of the organic phase for analysis of premnaspirodiene. The acetone and hexane used for extraction were each spiked with internal standards to aid in quantitation of the samples. The extracted samples were analyzed by GC and the amount of premnaspirodiene was calculated from the peak area.

[0109] Several mutant strains were identified that produced more or similar amounts of premnaspirodiene as the control strain ALX7-95 containing expressed HPS. They were given the designations ALX7-168.4, ALX7-168.10, ALX7-168.14, ALX7-168.19, ALX7-168.22, ALX7-168.23, ALX7-168.24, ALX7-168.25, and ALX7-183.69. The *ERG9* gene from these mutant strains was PCR amplified and the PCR product was sequenced to determine the mutations within each strain. A *HIS3*<sup>+</sup> strain of ALX7-168.25 was constructed and is designated ALX7-175.1.

### Example 3

#### Production of Premnaspirodiene in Fermentors

[0110] As described in Example 2, strain ALX7-175.1 was constructed for the production of premnaspirodiene. Production of premnaspirodiene in this strain was compared to that of strain ALX7-95 HPS, which is completely lacking in squalene synthase activity.

[0111] Production of premnaspirodiene was carried out in a 3-L fermentation tank (New Brunswick Bioflow 110). One liter of fermentation medium was prepared and autoclaved in the fermentation tank (20 g (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 20 g KH<sub>2</sub>PO<sub>4</sub>, 1 g NaCl, MgSO<sub>4</sub>·7H<sub>2</sub>O, 4 g yeast extract (Difco). Afterward the following components were added: 20 ml mineral solution (0.028% FeSO<sub>4</sub>·7H<sub>2</sub>O, 0.029% ZnSO<sub>4</sub>·7H<sub>2</sub>O, 0.008%

CuSO<sub>4</sub>·5H<sub>2</sub>O, 0.024% Na<sub>2</sub>MoO<sub>4</sub>·2H<sub>2</sub>O, 0.024% CoCl<sub>2</sub>·6H<sub>2</sub>O, 0.017% MnSO<sub>4</sub>·H<sub>2</sub>O, 1 mL HCl); 10 mL 50% glucose; 30 mL vitamin solution (0.001% biotin; 0.012% calcium pantothenate, 0.06% inositol, 0.012% pyridoxine-HCl, 0.012% thiamine-HCl); 10 mL 10% CaCl<sub>2</sub>, and 20 mL autoclaved soybean oil (purchased from local groceries). For ALX7-95 HPS, 1 mL of 50 mg/mL cholesterol in 100% ethanol was added.

[0112] The seed culture for inoculating the fermentation medium was prepared by inoculating 50 mL of SD medium for *ERG9* transformant strains. Non-transformant control cultures were grown in SDE medium (SD medium supplemented with 40 mg/L ergosterol). This culture was grown until early stationary phase (24-48 hr). One mL of this culture was inoculated into 500 mL of SD or SDE medium, as appropriate, and grown for 24 hr. A 50-mL aliquot (5% inoculum) was used to inoculate the one liter of medium.

[0113] The fermentor was maintained at 26° C. The air flow was 4.5 L/min and the dO<sub>2</sub> was maintained above 30% by adjusting the rpm. Furthermore, the pH was maintained at 4.5 using acetic acid and NaOH.

[0114] Once the glucose concentration was below 1 g/L, a feeding regimen was initiated such that the glucose in the fermentor was kept between 0 and 1 g/L. The glucose feed was made by mixing 1400 mL of 60% glucose and 328 mL of 12.5% yeast extract.

[0115] After 5 days, the air and agitation were turned off, and the oil was allowed to rise to the top of the tank and decanted.

#### Example 4

##### Comparison of Premnaspirodiene Production in *erg9* Mutants and *erg9::HIS3* Strains in Microbial Cultures

[0116] As described in Example 2, initial screening of mutants was conducted using microbial cultures. Mutants were further screened in microbial cultures by growing in media with or without 40 mg/L ergosterol supplementation and compared to ALX7-95 HPS grown with ergosterol (ALX7-95 HPS will not grow without ergosterol). Twenty-five isolates are compared in Figure 1. In Figure 1, the concentration of premnaspirodiene in mg/L is shown for each of the 25 isolates with and without

ergosterol supplementation. Several of these isolates produced more or comparable amounts of premnaspirodiene as the control culture ALX7-95 HPS. In general, these strains produced more premnaspirodiene in the absence of ergosterol than in its presence. Isolates 4, 10, 114, 19, 22, 23, 24, 25, and 69 were given strain designations ALX7-168.4, ALX7-168.10, ALX7-168.14, ALX7-168.19, ALX7-168.22, ALX7-168.23, ALX7-168.24, ALX7-168.25, and ALX7-83.169 respectively.

#### Example 5

##### Comparison of Premnaspirodiene Production in *erg9* Mutants and *erg9::HIS3* Strains in Fermentation Cultures

[0117] In this example, strains ALX7-95 HPS (*erg9::HIS3*) and ALX7-175.1, a histidine prototroph of ALX7-168.25 (*erg9<sup>def</sup>*) were grown in fermentors using the same protocol except for the presence or absence of cholesterol in the medium. At the end of the fermentation, premnaspirodiene was assayed by gas chromatography. As indicated in Table 1, the yields of premnaspirodiene in the fermentors were similar. However, because of faster growth and growth to higher density of the *erg9<sup>def</sup>* strain, more glucose was fed and consumed during the course of the fermentation. Because of the resultant higher volume, more total premnaspirodiene was produced by the *erg9<sup>def</sup>* culture grown under the same starting conditions.

Table 1

Strain	ERG9 Allele	Cell Density, OD <sub>600</sub>	Premnaspirodiene Titer, g/L	Final Volume, Liters	Total Yield, grams
ALX7-95 HPS	<i>erg9::HIS3</i>	48	3.9	1.1	4.3
ALX7-175.1	<i>erg9<sup>def</sup>25</i>	193	3.7	1.7	6.2

#### Example 6

##### Sequences of *erg9<sup>def</sup>* Mutants

[0118] The sequence of the wild type *ERG9* gene and sequences 245 base pairs upstream are shown in Figure 2. The *ERG9* alleles from strains described in Examples 2



and 4 were obtained by PCR amplification of genomic DNA from strains designations ALX7-168.4, ALX7-168.10, ALX7-168.14, ALX7-168.19, ALX7-168.22, ALX7-168.23, ALX7-168.24, ALX7-168.25, and ALX7-183.69. The resulting DNA was sequenced, and the sequences corresponding to those mutants are shown in Table 2. All mutant genes contain mutations in the *ERG9* coding region. Alleles 10, 23, 24, and 25 contain, in addition, mutations in the 245 base pair noncoding region of upstream of the gene.

Table 2

Mutant	Nucleotide position and change	Amino Acid position and change	Affect of change on amino acid
4	691 A→G	149 E→G	silent
	748 G→T	168 G→V	
	786 T→A	181 Y→N	
	1114 A→T	290 Q→L	
	1213 T→C	323 I→T	
	1290 T→C	349 L→L	
10	72 C→A	186 V→I	non-coding
	110 Δ A		non-coding
	801 G→A		
14	989 T→A	248 P→P	silent
	1112 G→A	289 E→E	silent
	1220 G→A	325 K→K	silent
	1233 T→C	330 Y→H	
19	786 T→A	181 Y→N	silent
	1025 A→G	260 Q→Q	
	1056 T→A	271 L→I	
	1068 A→G	275 S→G	
	1203 A→G	320 N→D	
22	886 T→C	214 M→T	
	969 A→G	242 I→V	
	1075 T→C	277 V→A	
	1114 A→T	290 Q→L	

23	84 T→A		non-coding
	283 A→T	13 E→V	
	424 T→C	60 L→P	
	440 A→G	65 R→R	silent
	1076 T→C	277 V→V	silent
24	619 A→T	125 D→V	
	634 T→C	130 L→P	
	962 C→T	239 P→P	silent
25	150 A→T		non-coding
	410 T→G	55 A→A	silent
	411 G→T	56 V→L	
	1248 T→C	335 S→P	
69	510 C→T	89 H→Y	
	573 T→C	110 F→L	
	918 A→G	224 R→G	
	997 A→G	251 K→G	

[0119] As used in this specification and in the appended claims, the singular forms include the plural forms. For example the terms “a,” “an,” and “the” include plural references unless the content clearly dictates otherwise. Additionally, the term “at least” preceding a series of elements is to be understood as referring to every element in the series. The inventions illustratively described herein can suitably be practiced in the absence of any element or elements, limitation or limitations, not specifically disclosed herein. Thus, for example, the terms “comprising,” “including,” “containing,” etc. shall be read expansively and without limitation. Additionally, the terms and expressions employed herein have been used as terms of description and not of limitation, and there is no intention in the use of such terms and expressions of excluding any equivalents of the future shown and described or any portion thereof, and it is recognized that various modifications are possible within the scope of the invention claimed.

[0120] Although the present invention has been described in considerable detail with reference to certain preferred embodiments, other embodiments are possible. The steps disclosed for the present methods, for example, are not intended to be limiting nor

are they intended to indicate that each step is necessarily essential to the method, but instead are exemplary steps only. Therefore, the scope of the appended claims should not be limited to the description of preferred embodiments contained in this disclosure. All references cited herein are incorporated by reference in their entirety.

We claim:

1. An isolated nucleic acid molecule that encodes a squalene synthase enzyme that, when present and expressed *in vivo* in a eukaryotic microbial host as the only squalene synthase species, catalyzes the synthesis of squalene at a sufficiently high rate that supplementation of the eukaryotic microbial host with a sterol is not required for growth and results in a reduced cellular squalene synthase activity.
2. The isolated nucleic acid molecule of claim 1 wherein the eukaryotic microbial host is a fungal microbial host.
3. The isolated nucleic acid molecule of claim 2 wherein the fungal microbial host is a yeast microbial host.
4. The isolated nucleic acid molecule of claim 3 wherein the yeast microbial host is selected from the group consisting of *Saccharomyces*, *Zygosaccharomyces*, *Kluyveromyces*, *Candida*, *Hansenula*, *Debaryomyces*, *Mucor*, *Pichia* and *Torulopsis*.
5. The isolated nucleic acid molecule of claim 4 wherein the yeast microbial host is *Saccharomyces cerevisiae*.
6. The isolated nucleic acid molecule of claim 1 wherein the squalene synthase enzyme encoded by the nucleic acid molecule has a reduced  $V_{\max}$  for squalene synthesis.
7. The isolated nucleic acid molecule of claim 1 wherein the squalene synthase enzyme encoded by the nucleic acid molecule has an increased Michaelis constant ( $K_m$ ) for its farnesyl pyrophosphate (FPP) substrate.

8. The isolated nucleic acid molecule of claim 1 that encodes a squalene synthase enzyme that, when expressed *in vivo* in the eukaryotic microbial host, produces squalene at a rate of less than 75% of the wild-type enzyme.

9. The isolated nucleic acid molecule of claim 8 that encodes a squalene synthase enzyme that, when expressed *in vivo* in the eukaryotic microbial host, produces squalene at a rate of less than 50% of the wild-type enzyme.

10. The isolated nucleic acid molecule of claim 9 that encodes a squalene synthase enzyme that, when expressed *in vivo* in the eukaryotic microbial host, produces squalene at a rate of less than 25% of the wild-type enzyme.

11. The isolated nucleic acid molecule of claim 1 wherein the isolated nucleic acid molecule has at least one change from a nucleic acid molecule that includes the coding region for the wild-type *ERG9* gene and its flanking sequences, including the sequences both upstream and downstream from the coding region, such that either: (a) the specific activity of the resulting squalene synthase enzyme is reduced through at least one change in the amino acid sequence of the enzyme expressed from the nucleic acid molecule; or (b) the *in vivo* activity of the enzyme is reduced through a reduction in transcription, a reduction in translation, or a reduction of enzyme stability.

12. The isolated nucleic acid molecule of claim 1 wherein the nucleic acid molecule, when expressed *in vivo*, causes a host cell to produce a greater concentration of an isoprenoid in grams of isoprenoid per liter of culture than a corresponding host cell expressing a wild-type nucleic acid molecule.

13. The isolated nucleic acid molecule of claim 5 wherein the nucleic acid molecule is selected from the group consisting of at least one of the mutants listed in Table 2.

14. The isolated nucleic acid molecule of claim 13 wherein one or more of the silent amino acid mutations in the mutant is omitted.

15. The isolated nucleic acid molecule of claim 13 in which the isolated nucleic acid molecule is an isolated nucleic acid molecule in which the squalene synthase enzyme encoded by the isolated nucleic acid molecule differs by one or more conservative amino acid substitutions.

16. An isolated nucleic acid molecule that is at least 95% identical to an isolated nucleic acid molecule selected from the group consisting of at least one of the mutants listed in Table 2.

17. The isolated nucleic acid molecule of claim 16 that is at least 97.5% identical to an isolated nucleic acid molecule selected from group consisting of at least one of the mutants listed in Table 2.

18. The isolated nucleic acid molecule of claim 17 that is at least 99% identical to an isolated nucleic acid molecule selected from the group consisting of at least one of the mutants listed in Table 2.

19. The isolated nucleic acid molecule of claim 18 that is at least 99.5% identical to an isolated nucleic acid molecule selected from the group consisting of at least one of the mutants listed in Table 2.

20. The isolated nucleic acid molecule of claim 19 that is at least 99.8% identical to an isolated nucleic acid molecule selected from the group consisting of at least one of the mutants listed in Table 2.

21. The isolated nucleic acid molecule of claim 1 wherein the nucleic acid molecule further comprises at least one additional nucleic acid sequence at the 5'-terminus, the 3'-terminus, or at both the 5'-terminus and 3'-terminus, to which the isolated nucleic acid molecule of claim 1 is operatively linked.

22. The isolated nucleic acid molecule of claim 21 wherein the at least one additional nucleic acid sequence is selected from the group consisting of linkers, adaptors, restriction endonuclease cleavage sites, regulatory sequences, and coding sequences.

23. A vector comprising the isolated nucleic acid molecule of claim 1.

24. The vector of claim 23 wherein the vector is capable of replication in a prokaryotic organism.

25. The vector of claim 23 wherein the vector is capable of replication in a eukaryotic organism.

26. The vector of claim 23 wherein the vector is capable of integration into the genome of a mammalian cell line.

27. The vector of claim 23 wherein the vector further comprises at least one additional nucleic acid sequence at the 5'-terminus, the 3'-terminus, or at both the 5'-terminus and 3'-terminus, to which the isolated nucleic acid molecule encoding a squalene synthase enzyme is operatively linked.

28. The vector of claim 27 wherein the at least one additional nucleic acid sequence is selected from the group consisting of linkers, adaptors, restriction endonuclease cleavage sites, regulatory sequences, and coding sequences.

29. A host cell including therein the nucleic acid of claim 1.

30. The host cell of claim 29 wherein the host cell is a prokaryotic cell.
31. The host cell of claim 29 wherein the host cell is a eukaryotic cell.
32. The host cell of claim 29 wherein the host cell further comprises at least one additional nucleic acid sequence at the 5'-terminus, the 3'-terminus, or at both the 5'-terminus and 3'-terminus, to which the isolated nucleic acid molecule encoding a squalene synthase enzyme is operatively linked.
33. The host cell of claim 29 wherein the at least one additional nucleic acid sequence is selected from the group consisting of linkers, adaptors, restriction endonuclease cleavage sites, regulatory sequences, and coding sequences.
34. A host cell transformed or transfected with the vector of claim 23.
35. A variant squalene synthase enzyme that, when present and expressed *in vivo* in a eukaryotic microbial host as the only squalene synthase species, catalyzes the synthesis of squalene at a sufficiently high rate that supplementation of the eukaryotic microbial host with a sterol is not required for growth and that results in reduced cellular squalene synthase activity.
36. The variant squalene synthase enzyme of claim 35 wherein the eukaryotic microbial host is a fungal microbial host.
37. The variant squalene synthase enzyme of claim 36 wherein the fungal microbial host is a yeast microbial host.



38. The variant squalene synthase enzyme of claim 37 wherein the yeast microbial host is selected from the group consisting of *Saccharomyces*, *Zygosaccharomyces*, *Kluyveromyces*, *Candida*, *Hansenula*, *Debaryomyces*, *Mucor*, *Pichia* and *Torulopsis*.

39. The variant squalene synthase enzyme of claim 38 wherein the *Saccharomyces* yeast microbial host is *Saccharomyces cerevisiae*.

40. The variant squalene synthase enzyme of claim 35 wherein the squalene synthase enzyme has a reduced  $V_{\max}$  for squalene synthesis.

41. The variant squalene synthase enzyme of claim 35 wherein the squalene synthase enzyme has an increased  $K_m$  for its FPP substrate.

42. The variant squalene synthase enzyme of claim 35 wherein the squalene synthase enzyme produces squalene *in vivo* in the eukaryotic microbial host at a rate of less than 75% of the wild-type enzyme.

43. The variant squalene synthase enzyme of claim 42 wherein the squalene synthase enzyme produces squalene *in vivo* in the eukaryotic microbial host at a rate of less than 50% of the wild-type enzyme.

44. The variant squalene synthase enzyme of claim 43 wherein the squalene synthase enzyme produces squalene *in vivo* in the eukaryotic microbial host at a rate of less than 25% of the wild-type enzyme.

45. The variant *S. cerevisiae* squalene synthase enzyme of claim 39 wherein the variant *S. cerevisiae* squalene synthase enzyme is selected from the group consisting of at least one of the mutants listed in Table 2.

46. The variant *S. cerevisiae* squalene synthase enzyme of claim 45 wherein the squalene synthase enzyme is a variant squalene synthase enzyme in which the squalene synthase enzyme differs by one or more conservative amino acid substitutions from a variant squalene synthase enzyme selected from the group consisting of at least one of the mutants listed in Table 2.

47. A method of isolating a defective *ERG9* gene comprising the steps of:

- (a) isolating a wild-type *ERG9* gene to produce an isolated wild-type *ERG9* gene;
- (b) subjecting the isolated wild-type *ERG9* gene to mutagenesis to generate a pool of *ERG9* mutants;
- (c) transforming mutants from the pool of *erg9* mutants generated in step (b) into a strain of a eukaryotic microbial host that contains an expressed terpene synthase gene that produces a detectable and measurable terpene product, the strain of the eukaryotic microbial host being transformed in such a manner that replacement of the preexisting *ERG9* allele with an *erg9* mutation allows the strain to grow in a sterol-free medium; and
- (d) isolating a transformant from step (c) that produces a level of terpene product at least equivalent to the level of terpene product produced by a strain of the eukaryotic microbial host expressing the terpene synthase gene that requires a sterol in the medium for growth.

48. The method of claim 47 wherein the step of isolating the wild-type *ERG9* gene is performed by a nucleic amplification method.

49. The method of claim 47 wherein the *ERG9* gene is a fungal *ERG9* gene.

50. The method of claim 49 wherein the fungal *ERG9* gene is a *Saccharomyces* *ERG9* gene.

51. The method of claim 50 wherein the *Saccharomyces ERG9* gene is a *Saccharomyces cerevisiae ERG9* gene.

52. The method of claim 47 wherein the mutagenesis is carried out by performing error-prone PCR.

53. The method of claim 52 wherein the error-prone PCR is performed using one or more DNA polymerase enzymes that have higher misinsertion and misextension rates than wild-type polymerase enzymes.

54. The method of claim 47 wherein the eukaryotic microbial host is a fungal microbial host.

55. The method of claim 54 wherein the fungal microbial host is a yeast microbial host.

56. The method of claim 55 wherein the yeast microbial host is a yeast of the genus *Saccharomyces*.

57. The method of claim 56 wherein the yeast of the genus *Saccharomyces* is *Saccharomyces cerevisiae*.

58. The method of claim 47 wherein the terpene synthase gene that produces a detectable and measurable terpene product is a *Hyoscyamus muticus* premnaspirodiene synthase (HPS) gene.

59. The method of claim 47 wherein the terpene synthase gene that produces a detectable and measurable terpene product is one that produces a product detectable and measurable by gas chromatography.

60. The method of claim 57 wherein the strain of *S. cerevisiae* that contains an expressed terpene synthase gene that produces a detectable and measurable terpene product is ALX7-95 (*his3, trp1, erg9::HIS3, HMGcat/TRP1::rDNA, dpp1*), a leucine prototroph of strain CALI-5, that contains a plasmid expressing the *Hyoscyamus muticus* premnaspirodiene synthase (HPS) gene.

61. A method of isolating a variant squalene synthase enzyme comprising the steps of:

(a) culturing a host cell that expresses the variant squalene synthase enzyme that, when present and expressed in vivo in a eukaryotic microbial host as the only squalene synthase species, catalyzes the synthesis of squalene at a sufficiently high rate that supplementation of the eukaryotic microbial host with a sterol is not required for growth and that results in reduced cellular squalene synthase activity; and

(b) isolating the variant squalene synthase enzyme from the host cell.

62. The method of claim 61 wherein the method further comprises the step of purifying the isolated variant squalene synthase enzyme.

63. The method of claim 61 wherein the variant squalene synthase enzyme is a variant *Saccharomyces cerevisiae* squalene synthase enzyme.

64. A method of producing an isoprenoid using a mutated *ERG9* gene, which mutated *ERG9* gene encodes the variant squalene synthase enzyme that, when present and expressed in vivo in a eukaryotic microbial host as the only squalene synthase species, catalyzes the synthesis of squalene at a sufficiently high rate that supplementation of the eukaryotic microbial host with a sterol is not required for growth and that results in reduced cellular squalene synthase activity, the method comprising the steps of:

(a) providing a host cell that includes a mutated *ERG9* gene that encodes the variant squalene synthase enzyme and at least one isoprenoid synthase gene;

(b) allowing the host cell to produce farnesyl pyrophosphate and to synthesize the isoprenoid from the farnesyl pyrophosphate; and

(c) isolating the isoprenoid synthesized by the host cell.

65. The method of claim 64 wherein the host cell is a *Saccharomyces cerevisiae* cell.

66. The method of claim 64 wherein the variant squalene synthase enzyme is a variant *Saccharomyces cerevisiae* squalene synthase enzyme.

67. The method of claim 64 wherein the isoprenoid synthase gene is selected from the group consisting of the *Hyoscyamus muticus* vetispiradiene synthase gene, the *H. muticus* premnaspirodiene synthase gene, the *Nicotiana tabacum* 5-epi-aristolochene synthase gene, a citrus valencene synthase gene, the *Capsicum annuum* 5-epi-aristolochene synthase gene, the *Mentha piperita* E- $\beta$ -farnesene synthase gene, the *Abies grandis*  $\delta$ -selenene synthase gene, the *Abies grandis*  $\gamma$ -humulene synthase gene, the *Abies grandis* - $\alpha$ -bisabolene synthase gene, the *Gossypium arboreum* 6-cadinene synthase gene, the *Lycopersicon esculentum* germacrene C synthase gene, the *Artemisia annua* epi-cedrol synthase gene, the *Artemisia annua* amorpho-4,11-diene synthase gene, the *Lactuca sativa* germacrene A synthase gene, the *Cichorium intybus* germacrene A synthase gene, and the *Solidago canadensis* germacrene A synthase gene, and mutants and protein engineered variants thereof.

68. The method of claim 67 wherein the isoprenoid synthase gene is the *H. muticus* premnaspirodiene synthase gene and the isoprenoid produced is premnaspirodiene.

69. The method of claim 68 wherein the premnaspirodiene is isolated by (i) sequestering the premnaspirodiene by binding it to a hydrophobic resin; and (ii) isolating the premnaspirodiene from the hydrophobic resin.

70. The method of claim 68 wherein the premnaspirodiene is isolated by using a two-phase system in which a non-polar solvent, substantially immiscible with an aqueous phase, is added to the fermentation broth, the nonpolar solvent is separated from the aqueous phase of the fermentation broth, and the premnaspirodiene is removed from the non-polar phase by distillation.

71. The method of claim 70 wherein the non-polar solvent is an oil.

72. The method of claim 64 further comprising the step of performing at least one additional reaction on the isolated isoprenoid in order to transform the isolated isoprenoid into another isoprenoid or related compound.

73. The method of claim 72 wherein the at least one additional reaction is selected from the group consisting of oxidation, hydroxylation, alkylation, and halogenation.

74. The method of claim 73 wherein the at least one additional reaction is an oxidation or hydroxylation and is carried out by use of a cytochrome P450 enzyme.

75. A method of producing an isoprenoid using a mutated *ERG9* gene, which mutated *ERG9* gene encodes the variant squalene synthase enzyme that, when present and expressed in vivo in a eukaryotic microbial host as the only squalene synthase species, catalyzes the synthesis of squalene at a sufficiently high rate that supplementation of the eukaryotic microbial host with a sterol is not required for growth and that results in reduced cellular squalene synthase activity, the method comprising the steps of:

- (a) providing a host cell that includes a mutated *ERG9* gene that encodes the variant squalene synthase enzyme and at least one isoprenoid synthase gene;
- (b) allowing the host cell to produce farnesyl pyrophosphate;
- (c) isolating farnesyl pyrophosphate from the host cell;

(d) reacting the farnesyl pyrophosphate *in vitro* with one or more isoprenoid synthases to synthesize the isoprenoid; and

(e) isolating the isoprenoid.

76. The method of claim 75 wherein the host cell is a *Saccharomyces cerevisiae* cell.

77. The method of claim 76 wherein the variant squalene synthase enzyme is a variant *Saccharomyces cerevisiae* squalene synthase enzyme.

78. The method of claim 76 wherein the isoprenoid synthase is selected from the group consisting of the *Hyoscyamus muticus* vetispiradiene synthase, the *H. muticus* premnaspirodiene synthase, the *Nicotiana tabacum* 5-epi-aristolochene synthase, a citrus valencene gene, the *Capsicum annuum* 5-epi-aristolochene synthase, the *Mentha piperita* E- $\beta$ -farnesene synthase, the *Abies grandis*  $\delta$ -selenene synthase, the *Abies grandis*  $\gamma$ -humulene synthase, the *Abies grandis* - $\alpha$ -bisabolene synthase, the *Gossypium arboreum* 6-cadinene synthase, the *Lycopersicon esculentum* germacrene C synthase, the *Artemisia annua* epi-cedrol synthase, the *Artemisia annua* amorpho-4,11-diene synthase, the *Lactuca sativa* germacrene A synthase, the *Cichorium intybus* germacrene A synthase, and the *Solidago canadensis* germacrene A synthase, and mutants and protein engineered variants thereof.

79. The method of claim 78 wherein the isoprenoid synthase the *H. muticus* premnaspirodiene synthase and the isoprenoid produced is premnaspirodiene.

80. The method of claim 79 wherein the premnaspirodiene is isolated by (i) sequestering the premnaspirodiene by binding it to a hydrophobic resin; (ii) and isolating the premnaspirodiene from the hydrophobic resin.

81. The method of claim 79 wherein the premnaspirodiene is isolated by using a two-phase system in which a non-polar solvent, substantially immiscible with an aqueous phase, is added to the fermentation broth, the nonpolar solvent is separated from the aqueous phase of the fermentation broth, and the premnaspirodiene is removed from the non-polar phase by distillation.

82. The method of claim 81 wherein the non-polar solvent is an oil.

83. The method of claim 77 further comprising the step of performing at least one additional reaction on the isolated isoprenoid in order to transform the isolated isoprenoid into another isoprenoid or related compound.

84. The method of claim 83 wherein the at least one additional reaction is selected from the group consisting of oxidation, hydroxylation, alkylation, and halogenation.

85. The method of claim 84 wherein the at least one additional reaction is an oxidation or hydroxylation and is carried out by use of a cytochrome P450 enzyme.



Figure 1




Figure 2

10 20 30 40 50  
GCCCATCTTCAACAACAATACCGACTTACCATCCTATTTGCTTTGCCCTTTT  
7-162.1  
60 70 80 90 100  
TCTTTTCCACTGCACTTTGCAATCGGAAGGCGTTATCGGTTTTGGGTTTAGTG  
110 120 130 140 150  
CCTAAACGAGCAGCGAGAACACGACCACGGGCTATATAAATGGAAAGTTAGG  
160 170 180 190 200  
ACAGGGGCAAAGAATAAGAGCACAGAAGAAGAGAAAAGACGAAGAGCAGAAG  
210 220 230 240 250 260  
CGGAAAACGTATACACGTCACATATCACACACACAATGGGAAAGCTATTA  
M G K L L>  
270 280 290 300 310  
CAATTGGCATTGCATCCGGTCGAGATGAAGGCAGCTTTGAAGCTGAAGTTTT  
Q L A L H P V E M K A A L K L K F>  
320 330 340 350 360  
GCAGAACACCGCTATTCTCCATCTATGATCAGTCCACGTCTCCATATCTCTT  
C R T P L F S I Y D Q S T S P Y L L  
370 380 390 400 410  
GCACTGTTTTCGAACTGTTGAACCTTGACCTCCAGATCGTTTGCTGCTGTGATC  
H C F E L L N L T S R S F A A V I>  
420 430 440 450 460  
AGAGAGCTGCATCCAGAATTGAGAACTGTGTTACTCTCTTTTATTTGATTT  
R E L H P E L R N C V T L F Y L I>  
470 480 490 500 510 520  
TAAGGGCTTTGGATACCATCGAAGACGATATGTCCATCGAACACGATTTGAA  
L R A L D T I E D D M S I E H D L K>  
530 540 550 560 570  
AATTGACTTGTTGCGTCACTTCCACGAGAAATTGTTGTTAACTAAATGGAGT  
I D L L R H F H E K L L L T K W S>

Figure 2 cont.

580 590 600 610 620  
TTCGACGGAAATGCCCCCGATGTGAAGGACAGAGCCGTTTGTGACAGATTTTCG  
F D G N A P D V K D R A V L T D F>

630 640 650 660 670  
AATCGATTCTTATTGAATTCCACAAATTGAAACCAGAATATCAAGAAGTCAT  
E S I L I E F H K L K P E Y Q E V I>

680 690 700 710 720  
CAAGGAGATCACCGAGAAAATGGGTAAATGGTATGGCCGACTACATCTTAGAT  
K E I T E K M G N G M A D Y I L D>

730 740 750 760 770 780  
GAAAATTACAACCTTGAATGGGTTGCAAACCGTCCACGACTACGACGTGTACT  
E N Y N L N G L Q T V H D Y D V Y>

790 800 810 820 830  
GTCACTACGTAGCTGGTTTGGTTCGGTGATGGTTTGAACCGTTTGTATTGTCAT  
C H Y V A G L V G D G L T R L I V I>

840 850 860 870 880  
TGCCAAGTTTGCCAACGAATCTTTGTATTCTAATGAGCAATTGTATGAAAGC  
A K F A N E S L Y S N E Q L Y E S>

890 900 910 920 930  
ATGGGTCTTTTCCTACAAAAAACCAACATCATCAGAGATTACAATGAAGATT  
M G L F L Q K T N I I R D Y N E D>

940 950 960 970 980  
TGGTCGATGGTAGATCCTTCTGGCCCAAGGAAATCTGGTCACAATACGCTCC  
L V D G R S F W P K E I W S Q Y A P>

990 1000 1010 1020 1030 1040  
TCAGTTGAAGGACTTCATGAAACCTGAAAACGAACAACCTGGGGTTGGACTGT  
Q L K D F M K P E N E Q L G L D C>

1050 1060 1070 1080 1090  
ATAAACACCTCGTCTTAAACGCATTGAGTCATGTTATCGATGTGTTGACTT  
I N H L V L N A L S H V I D V L T>

Figure 2 cont.

=

1100 1110 1120 1130 1140  
ATTTGGCCGGTATCCACGAGCAATCCACTTTCCAATTTTGTGCCATTCCCCA  
Y L A G I H E Q S T F Q F C A I P Q>

1150 1160 1170 1180 1190  
AGTTATGGCCATTGCAACCTTGGCTTTGGTATTCAACAACCGTGAAGTGCTA  
V M A I A T L A L V F N N R E V L>

1200 1210 1220 1230 1240  
CATGGCAATGTAAAGATTCGTAAGGGTACTACCTGCTATTTAATTTTGAAAT  
H G N V K I R K G T T C Y L I L K>

1250 1260 1270 1280 1290 1300  
CAAGGACTTTGCGTGGCTGTGTCGAGATTTTTGACTATTACTTACGTGATAT  
S R T L R G C V E I F D Y Y L R D I>

1310 1320 1330 1340 1350  
CAAATCTAAATTGGCTGTGCAAGATCCAAATTTCTTAAATTTGAACATTCAA  
K S K L A V Q D P N F L K L N I Q>

1360 1370 1380 1390 1400  
ATCTCCAAGATCGAACAGTTTATGGAAGAAATGTACCAGGATAAATTACCTC  
I S K I E Q F M E E M Y Q D K L P>

1410 1420 1430 1440 1450  
CTAACGTGAAGCCAAATGAAACTCCAATTTTCTTGAAAGTTAAAGAAAGATC  
P N V K P N E T P I F L K V K E R S>

1460 1470 1480 1490 1500  
CAGATACGATGATGAATTGGTTCCAACCCAACAAGAAGAAGAGTACAAGTTC  
R Y D D E L V P T Q Q E E E Y K F>

1510 1520 1530 1540 1550 1560  
AATATGGTTTTATCTATCATCTTGTCCGTTCTTCTTGGGTTTTATTATATAT  
N M V L S I I L S V L L G F Y Y I>

1570 1580 1590 1600 1610  
ACACTTTACACAGAGCGTGAAGTCTGCGCCAATAACATAAACAAACAACTC

Figure 2 cont.

Y T L H R A \*

1620 1630

CGAACAATAACTAAGTACT

7-162.2